

Supplementary Information

Title: Predictive Value of Waist Circumference and Body Mass Index for Lifestyle-Related Diseases in an Aging Asian Population

Contents

Figure S1. Violin Plots of BMI and Waist Circumference in Male and Female Participants

Figure S2. Violin Plots of Triglycerides, Total Cholesterol, HDL-C, and LDL-C in Male and Female Participants

Figure S3. Violin Plot of Fasting Blood Glucose Distribution in Male and Female Participants

Figure S4 ROC Curves for Waist Circumference and Body Mass Index in Predicting Type 2 Diabetes Mellitus

Figure S5. ROC Curves for Waist Circumference and Body Mass Index in Predicting Dyslipidemia

Figure S1. Violin Plots of BMI and Waist Circumference in Male and Female Participants




Figure S2. Violin Plots of Triglycerides, Total Cholesterol, HDL-C, and LDL-C in Male and Female Participants




Figure S3. Violin Plot of Fasting Blood Glucose Distribution in Male and Female Participants






Figure S4. ROC Curves for Waist Circumference and Body Mass Index in Predicting Type 2 Diabetes Mellitus



Panels A, B, and C show the predictive performance of waist circumference for T2DM in all individuals, males, and females, respectively; Panels D, E, and F show the predictive performance of body mass index for T2DM in all individuals, males, and females, respectively.

Figure S5. ROC Curves for Waist Circumference and Body Mass Index in Predicting Dyslipidemia



Panels A, B, and C show the predictive performance of waist circumference for dyslipidemia in all individuals, males, and females, respectively. Panels D, E, and F show the predictive performance of body mass index for dyslipidemia in all individuals, males, and females, respectively.