

Type of Article: Original Research

Title: Comparative Efficacy and Safety of Semaglutide, Cagrilintide, and CagriSema in Adults with Overweight or Obesity: A Bayesian Network Meta-Analysis

Authors: Yange Meng¹, Ting Jin², Luping Ren³, Michelle L Thompson⁴, Yue Wang⁵

¹ Department of General Practice, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China

² Department of Endocrinology, Sir Run Run Shaw Hospital, Zhejiang University School of Medicine, Hangzhou, China

³ Endocrinology Department, Hebei General Hospital, Shijiazhuang, China

⁴ Lifestyle Medicine, University of Pittsburgh Medical Center, Pittsburgh, USA

⁵ UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, USA

Correspondence: Yue Wang

5115 Centre Ave, Pittsburgh, PA 15232, United States

UPMC Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA, 15213, U.S.A.

E-mail: yue.wang@pitt.edu

Tel: 412-641-4530

19

Clinical Question Box

20 In adults with overweight or obesity, does the combination of Semaglutide and Cagrilintide
21 provide the most effective weight loss while maintaining an acceptable safety profile?

22 Evidence suggests that the combination therapy, known as CagriSema, produces the greatest
23 reductions in both body weight and waist circumference. Semaglutide alone results in substantial
24 weight loss with additional glycemic benefits, whereas Cagrilintide is better tolerated but less
25 effective. Liraglutide appears both less effective and less tolerable. None of these agents are
26 associated with an increased risk of serious adverse events, although gastrointestinal side effects
27 are more frequent with GLP-1–based therapies.

28

Introduction: Incretin-based therapies, including glucagon-like peptide-1 (GLP-1) receptor agonists and amylin analogues, have emerged as promising treatments for obesity. However, the relative efficacy and safety of Semaglutide, Cagrilintide, Liraglutide, and the novel combination therapy CagriSema remain uncertain.

Methods: A systematic review and network meta-analysis of randomized controlled trials was conducted in adults with overweight or obesity (BMI 25–40 kg/m²), with or without comorbidities. Outcomes included reductions in body weight and waist circumference, glycated hemoglobin (HbA1c), serious adverse events (SAEs), overall treatment discontinuation, and discontinuation due to gastrointestinal symptoms. Mean differences (MDs) and log odds ratios (LORs) with 95% confidence intervals (CIs) were calculated.

Results: Twelve trials involving 25,401 patients were included, with follow-up periods ranging from 20 to 104 weeks. CagriSema achieved the greatest reduction in body weight (MD: 17.7%, 95% CI: 14.2–21.3), followed by Semaglutide (MD: 11.1%, 95% CI: 9.5–12.7) and Cagrilintide (MD: 5.9%, 95% CI: 1.5–10.0). CagriSema also showed the largest reduction in waist circumference (MD: –13.4 cm, 95% CI: –17.1 to –9.7), followed by Semaglutide (MD: –8.4 cm, 95% CI: –10.0 to –7.0) and Cagrilintide (MD: –3.6 cm, 95% CI: –7.0 to 0.0). None of the treatments increased the risk of SAEs compared with placebo. However, Semaglutide was associated with higher rates of treatment discontinuation (LOR: 0.6, 95% CI: 0.1–1.2). Gastrointestinal-related discontinuation was highest with Liraglutide (LOR: 3.6, 95% CI: 1.5–5.6), followed by CagriSema (LOR: 2.5, 95% CI: 1.3–3.7) and Semaglutide (LOR: 1.7, 95% CI: 1.0–2.3); no significant effect was observed with Cagrilintide.

Conclusions: CagriSema provides the greatest reductions in weight and waist circumference. Semaglutide combines substantial weight loss with glycemic benefit, while Cagrilintide is better tolerated but less effective. Meanwhile, Liraglutide is both less effective and less tolerable.

Keywords: Semaglutide, Cagrilintide, CagriSema, obesity, overweight, network meta-analysis

Just Accepted

Introduction

Obesity has become a defining health challenge of the modern era. According to the World Health Organization (WHO), more than 2.5 billion adults worldwide are overweight, and nearly 900 million meet criteria for obesity.¹ These conditions substantially increase the risk of cardiometabolic disease, nonalcoholic fatty liver disease, sleep-disordered breathing, and certain malignancies, and are associated with diminished life expectancy.² Beyond their clinical consequences, obesity and overweight impose a substantial societal burden, with far-reaching psychological, social, and economic costs.³ Despite the central role of lifestyle modification, long-term adherence to dietary and behavioral interventions is difficult, and durable, clinically meaningful weight loss is uncommon.⁴ As a result, pharmacologic therapy has become a critical component of comprehensive obesity management.⁵

Among the available agents, glucagon-like peptide-1 receptor agonists (GLP-1 RAs) have redefined therapeutic expectations.⁶ Semaglutide, a long-acting GLP-1 RA, has consistently produced weight loss of unprecedented magnitude in randomized controlled trials (RCTs).⁷ In the STEP program, once-weekly subcutaneous Semaglutide (2.4 mg) resulted in weight reductions approaching those typically seen after bariatric surgery, alongside broad improvements in cardiometabolic risk factors.⁸ The drug's safety profile is generally acceptable, though gastrointestinal adverse events, particularly nausea, vomiting, and diarrhea, are frequent.⁹ The striking efficacy of Semaglutide has intensified efforts to develop novel peptide-based therapies that target complementary pathways involved in appetite and energy balance. Liraglutide, an earlier GLP-1 RA, demonstrated proof of concept that incretin-based therapy can facilitate weight reduction, though with modest effects compared with Semaglutide.

One such complementary pathway involves amylin, a pancreatic hormone co-secreted with insulin that slows gastric emptying and promotes satiety.¹⁰ Cagrilintide, a next-generation, long-acting amylin analogue, has demonstrated reductions in food intake and clinically meaningful weight loss in both preclinical and early-phase clinical trials.¹¹ Notably, Cagrilintide has also shown additive benefits when combined with GLP-1 RA, offering a strong rationale for dual-hormone approaches.^{12,13} The fixed-dose combination of Cagrilintide and Semaglutide, known as CagriSema, represents a novel therapeutic approach designed to exploit this synergy.¹⁴ Early trials suggest that CagriSema may achieve greater weight loss than either agent alone, with parallel improvements in metabolic outcomes and an acceptable safety profile.¹⁵ These encouraging findings have fueled considerable interest in dual-target therapies to address the limitations of monotherapy and meet the unmet needs of patients with obesity. However, direct head-to-head comparisons of Semaglutide, Cagrilintide, and CagriSema remain limited.

Bayesian network meta-analysis (NMA) provides a robust framework for evaluating such interventions. By synthesizing direct and indirect evidence, NMA allows simultaneous comparison of multiple agents and generates probabilistic treatment rankings to inform clinical decision-making. Given the rapidly evolving pharmacologic landscape, a comprehensive analysis of Semaglutide, Cagrilintide, and CagriSema is timely. The present study was designed to assess the relative efficacy and safety of these agents and to establish their comparative ranking in the treatment of overweight and obesity.

98 **Methods**

99 *Protocol and Registration*

100 This systematic review and NMA were conducted in accordance with the Preferred Reporting
101 Items for Systematic reviews and Meta-Analyses (PRISMA) 2020 and the PRISMA extension for
102 NMAs (PRISMA-NMA). The protocol was prospectively registered in the Open Science
103 Framework to ensure methodological transparency and reduce duplication (Registration ID:
104 osf.io/m3f6w).

105 *Eligibility Criteria*

106 Eligible studies were RCTs enrolling adults (≥ 18 years) with overweight (BMI ≥ 25 kg/m²) or
107 obesity (BMI ≥ 30 kg/m²). Interventions of interest included GLP-1 RAs (Liraglutide or
108 Semaglutide), Cagrilintide, or their combination, compared with placebo or active comparators.
109 All interventions were administered subcutaneously with a minimum follow-up of 20 weeks.
110 Studies including participants with prediabetes were eligible, whereas trials primarily focused on
111 type 2 diabetes were excluded. Open-label, crossover, or observational studies, as well as those
112 combining pharmacologic therapy with additional nonpharmacologic interventions, were excluded.

113 *Search Strategy*

114 A systematic search of PubMed, Embase, the Cochrane Library, and Web of Science was
115 performed from database inception through August 15, 2025. The search strategy combined
116 Medical Subject Headings (MeSH) and relevant keywords. Terms describing the study population
117 included “overweight” and “obesity.” Intervention terms included “semaglutide”, “cagrilintide,”
118 “CagriSema,” “GLP-1 receptor agonist,” and “amylin analogue.” Comparator terms included
119 “placebo” and “controlled.” No language restrictions were applied. Reference lists of eligible

studies and relevant systematic reviews were also screened. The detailed search strategy is provided in Table S1.

Study Selection

Titles and abstracts were independently screened by two reviewers, followed by full-text evaluations of potentially eligible reports. Disagreements were resolved through consensus or, when necessary, consultation with a third reviewer. The selection process was summarized in a PRISMA flow diagram.

Data Extraction

Data were extracted independently by two reviewers using a standardized electronic form. Extracted information included study characteristics (author, year, country, design, sample size, follow-up duration), participant characteristics (age, sex, baseline body mass index (BMI)), intervention details (drug, dose, route, schedule), and outcomes. When numerical data were unavailable, corresponding authors were contacted or data were estimated from published figures using validated extraction methods.

Risk of Bias Assessment

The quality of the included studies was assessed using the Cochrane Risk of Bias 2.0 tool, which evaluates randomization, deviations from intended interventions, missing outcome data, outcome measurement, and selective reporting. Each trial was rated as “low risk,” “some concerns,” or “high risk.” Assessments were performed independently by two reviewers, with discrepancies resolved through discussion.

Statistical Analysis

A Bayesian NMA was performed to compare the efficacy and safety of Semaglutide, Cagrilintide, and CagriSema. Random-effects models were used to account for heterogeneity. Analyses were conducted using Markov Chain Monte Carlo (MCMC) simulations in R with the gemtc and BUGSnet packages. Four chains of 50,000 iterations were run, following a burn-in of 20,000 iterations and a thinning interval of 10. Convergence was assessed by inspecting trace plots and the Gelman–Rubin statistic. Results were expressed as mean differences (MDs) or odds ratios (ORs) with 95% credible intervals (CIs).

Outcomes

The primary outcome was the percentage change in body weight from baseline. Secondary outcomes included changes in waist circumference, HbA1c, and safety outcomes such as serious adverse events (SAEs), discontinuations due to adverse events, and discontinuations due to gastrointestinal symptoms. Comparative effectiveness was further evaluated using the surface under the cumulative ranking curve (SUCRA), which estimates the probability of each treatment being the most effective.

Assessment of Inconsistency and Heterogeneity

Global inconsistency was evaluated using the design-by-treatment interaction model, while local inconsistency was examined with node-splitting analyses. Statistical heterogeneity was quantified using the posterior distribution of τ^2 , representing between-study variance. Publication bias was identified by Egger’s test.

Certainty of Evidence

The certainty of evidence was evaluated using the Grading of Recommendations Assessment, Development, and Evaluation (GRADE) framework adapted for NMAs. Considerations included

163 risk of bias, inconsistency, indirectness, imprecision, and publication bias. Evidence for each major
164 outcome was rated as high, moderate, low, or very low.

165

Just Accepted

Results

Study Selection and Characteristics

A total of 4,648 records were identified from four databases. After removing 216 duplicates, 4,432 records were screened. Of these, 4,311 records were excluded, leaving 121 reports, which were assessed for eligibility. A further 109 reports were excluded (reviews, non-RCTs, duplicates, oral admissions, or different backgrounds). Ultimately, 12 studies met the eligibility criteria (Figure S1). The included trials recruited adults with overweight or obesity (BMI 25–40), often with comorbidities such as osteoarthritis, prediabetes, or cardiovascular disease (Table 1).^{16–27} Most studies compared Semaglutide against placebo, while several also tested CagriSema, Cagrilintide, or Liraglutide. Sample sizes ranged from fewer than 40 participants in early-phase work to more than 17,000 in the largest cardiovascular outcomes trial. Follow-up durations spanned from 20 to 104 weeks. Participants were typically in their forties and fifties, though the cardiovascular trial enrolled an older cohort with a mean age of 62 years. Women comprised two-thirds to four-fifths of most study populations. Average BMI values generally ranged from 32 to 40 kg/m². Mean baseline HbA1c values were mostly close to the non-diabetic range (5.3–6.4%). All data included in the analysis were derived from studies using the approved doses of Cagrilintide or Semaglutide 2.4 mg.

Body Weight Reduction

The network of included studies assessing body weight is shown in Figure 1A. Eight trials evaluated Semaglutide versus placebo, with additional comparisons of Liraglutide versus placebo, Cagrilintide versus placebo, and CagriSema versus placebo. The estimated effects relative to placebo are presented in Figure 2A. CagriSema produced the greatest reduction in body weight

(MD: 17.7% [95% CI: 14.2–21.3]), followed by Semaglutide (11.9% [95% CI: 9.5–12.7]), Cagrilintide (5.9% [95% CI: 1.5–10.0]), and Liraglutide (4.3% [95% CI: 0.7–7.6]). The NMAs are summarized in Table 2. CagriSema was superior to all active comparators, compared with Semaglutide (MD: 6.6% [95% CI: 3.0–10.3]), Cagrilintide (MD: 11.8% [95% CI: 6.5–17.5]), and Liraglutide (MD: 13.4% [95% CI: 8.8–18.5]). Semaglutide was also superior to Cagrilintide (MD: 5.2% [95% CI: 1.0–9.7]) and Liraglutide (MD: 6.8% [95% CI: 3.5–10.4]). In contrast, Cagrilintide and Liraglutide did not differ significantly from each other (MD: 1.6% [95% CI: –2.6–5.8]).

SUCRA rankings (Figure S2) were as follows: CagriSema (0.999), Semaglutide (0.747), Cagrilintide (0.450), Liraglutide (0.298), and placebo (0.005). Egger’s test indicated no evidence of small-study effects ($p = 0.34$). Consistency assessment (Figure S3) showed no statistically significant differences ($p \geq 0.05$) between direct, indirect, and network estimates, indicating no notable inconsistency.

Waist Circumference Reduction

The network of included studies assessing waist circumference is shown in Figure 1B. Seven trials evaluated Semaglutide versus placebo, with additional comparisons of Liraglutide versus placebo, Cagrilintide versus placebo, and CagriSema versus placebo. The estimated effects relative to placebo are presented in Figure 2B. CagriSema produced the greatest reduction in waist circumference (MD: 13.4 cm [95% CI: 9.7–17.1]), followed by Semaglutide (8.4 cm [95% CI: 7.0–10.0]), Cagrilintide (3.6 cm [95% CI: 0.7–7.0]), and Liraglutide (2.3 cm [95% CI: –0.9–4.9]). The NMA results are summarized in Table 2. CagriSema was superior to all active comparators: Semaglutide (MD: 4.9 cm [95% CI: 0.9–8.9]), Cagrilintide (MD: 9.7 cm [95% CI: 4.8–15.0]), and Liraglutide (MD: 11.1 cm [95% CI: 6.7–16.0]). Semaglutide was also superior to Cagrilintide (MD: 4.8 cm [95% CI: 1.4–8.7]) and Liraglutide (MD: 6.2 cm [95% CI: 3.5–9.4]). In contrast,

Cagrilintide and Liraglutide did not differ significantly from each other (MD: 1.4 cm [95% CI: -1.9–5.0]).

SUCRA rankings (Figure S4) were as follows: CagriSema (0.996) ranked highest, followed by Semaglutide (0.751), Cagrilintide (0.453), Liraglutide (0.278), and placebo (0.022). Egger's test indicated no evidence of small-study effects ($p = 0.52$). Consistency assessment (Figure S5) showed no statistically significant differences ($p \geq 0.05$) between direct, indirect, and network estimates, indicating no notable inconsistency.

Serious Adverse Events (SAEs)

The network of included studies assessing SAEs is shown in Figure 1C. Ten trials evaluated Semaglutide versus placebo, two trials compared Liraglutide or Cagrilintide with placebo, and two trials compared CagriSema with Semaglutide. The estimated effects relative to placebo are presented in Figure 2C: Semaglutide (OR: 1.0 [95% CI: 0.8–1.3]), Cagrilintide (OR: 1.5 [95% CI: 0.8–2.8]), Liraglutide (OR: 1.6 [95% CI: 0.7–3.4]), and CagriSema (OR: 1.7 [95% CI: 0.9–3.0]). None of these agents demonstrated a statistically significant increase in the risk of SAEs compared with placebo.

The NMA results of log ORs are summarized in Table 2, showing that no treatment was superior or inferior to the others. SUCRA rankings (Figure S6) indicated that Semaglutide (0.812) and placebo (0.804) had the highest probabilities of being safest, followed by Cagrilintide (0.358), Liraglutide (0.318), and CagriSema (0.208). Egger's test showed no evidence of small-study effects ($p = 0.55$). Consistency assessment (Figure S7) revealed no statistically significant differences ($p \geq 0.05$) between direct, indirect, and network estimates, indicating no notable inconsistency.

233 *Treatment Discontinuation*

234 The network of included studies assessing treatment discontinuation was the same as for
235 SAEs (Figure 1C). The estimated effects relative to placebo are presented in Figure 2D.
236 Semaglutide (OR: 1.9 [95% CI: 1.1–3.5]) was associated with a significantly higher risk of
237 treatment discontinuation compared with placebo. In contrast, Cagrilintide (OR: 1.2 [95% CI: 0.4–
238 3.7]), Liraglutide (OR: 2.2 [95% CI: 0.7–6.6]), and CagriSema (OR: 2.4 [95% CI: 0.7–9.5]) did
239 not show statistically significant differences.

240 The NMA results of log ORs are summarized in Table 2. No treatment was superior or
241 inferior to the others, except for Semaglutide, which showed a higher risk of discontinuation than
242 placebo. SUCRA rankings (Figure S8) placed placebo highest (0.873), followed by Cagrilintide
243 (0.709), Semaglutide (0.373), Liraglutide (0.314), and CagriSema (0.232). Egger's test showed no
244 evidence of small-study effects ($p = 0.21$). Consistency assessment (Figure S9) revealed some
245 inconsistency in studies comparing Semaglutide or Cagrilintide with Liraglutide.

246 *Gastrointestinal Symptoms*

247 The network of included studies assessing treatment discontinuation due to gastrointestinal
248 symptoms is shown in Figure 1D. Seven trials evaluated Semaglutide versus placebo, and two
249 trials compared CagriSema with Semaglutide. The estimated effects relative to placebo are
250 presented in Figure 2E. Cagrilintide showed the lowest risk of treatment discontinuation (OR: 3.8,
251 95%CI: 0.68, 17.0) without significant statistical differences. Other agents showed a higher risk
252 of treatment discontinuation with Semaglutide (OR: 5.4, 95%CI: 2.7 to 10.0), CagriSema (OR:
253 12.0, 95%CI: 3.8, 41.0), and Liraglutide (OR: 28.0, 95%CI: 4.3, 270).

The NMA results of log ORs are summarized in Table 2. No treatment was superior or inferior to the others, except for Semaglutide, which showed a higher risk of discontinuation than placebo. SUCRA rankings (Figure S10) placed placebo at the highest for the lowest risk of discontinuation due to GS (0.985), followed by Cagrilintide (0.664), Semaglutide (0.556), CagriSema (0.225), and Liraglutide (0.070). Egger's test showed no evidence of small-study effects ($p = 0.21$). Consistency assessment (Figure S11) revealed no statistically significant differences ($p \geq 0.05$) between direct, indirect, and network estimates, indicating no notable inconsistency.

HbA1c

The network of included studies assessing waist circumference is shown in Figure S12. Four trials evaluated Semaglutide versus placebo, with additional comparisons of Liraglutide and Cagrilintide versus placebo. The estimated effects relative to placebo are presented in Figure 2F. Semaglutide produced the greatest reduction in waist circumference (MD: 0.45% [95% CI: 0.20–0.70]), followed by Liraglutide (MD: 0.22% [95% CI: –0.18–0.64]), and Cagrilintide (MD: 0.02% [95% CI: –0.49–54]). The NMA results are summarized in Table S2. Semaglutide was superior to the placebo (LOR: 0.4 [95%CI: 0.2, 0.7]), while no other agents were superior or inferior to the others.

SUCRA rankings (Figure S13) were as follows: Semaglutide (0.949) ranked highest, followed by Liraglutide (0.611), Cagrilintide (0.253), and placebo (0.187). Egger's test indicated no evidence of small-study effects ($p = 0.52$). Consistency assessment (Figure S14) showed no statistically significant differences ($p \geq 0.05$) between direct, indirect, and network estimates, indicating no notable inconsistency.

276 *Certainty of Evidence*

277 The risk of bias is presented in Figure S15. Two studies showed some concerns regarding
278 the overall risk of bias, while the remaining ten studies were assessed as having a low risk of bias.
279 Certainty of evidence is summarized in Table 3: high for reductions in body weight and waist
280 circumference, consistently showing that CagriSema was most effective, followed by Semaglutide.
281 Evidence for HbA1c reduction was also rated high for Semaglutide, but low to moderate for
282 Liraglutide and Cagrilintide, both of which showed no significant effect. For safety outcomes, the
283 certainty of evidence was moderate: no agent increased the risk of serious adverse events, although
284 Semaglutide and Liraglutide were associated with higher rates of treatment discontinuation,
285 particularly due to GS. Overall, the evidence supports a high level of confidence in efficacy
286 outcomes and a moderate level of confidence in safety and tolerability outcomes.

287

Discussion

This NMA consolidates evidence from randomized trials comparing CagriSema, Semaglutide, Cagrilintide, Liraglutide, and placebo in adults with overweight or obesity. The principal finding is that dual-agonist therapy with CagriSema provides the greatest reductions in body weight and waist circumference. Semaglutide also yields substantial benefits, achieving weight loss exceeding 10% alongside meaningful improvements in glycemic control. Liraglutide and Cagrilintide, while superior to placebo, produce comparatively more modest effects, generally achieving weight loss of more than 5%.²⁸ This NMA represents the ranking of all available agents across multiple outcomes in a single comparative framework, confirming the superiority of CagriSema and clarifying the trade-off between efficacy and tolerability. These results align with recent data from the REDEFINE program, which showed CagriSema achieving greater weight reductions than Semaglutide alone. This is consistent with earlier findings from the STEP program, confirming the superiority of Semaglutide over Liraglutide in weight management.^{29,30}

Different agents demonstrated distinct profiles when considering weight, glycemic outcomes, and safety. CagriSema consistently ranked highest for weight and waist circumference reduction, confirming its additive benefit over single-agent therapies and mirroring results from REDEFINE-1, where weight loss in adherent participants exceeded 20%.¹⁹ Semaglutide maintains a central role due to its dual impact on weight and glycemic control, supported by both trial and real-world data demonstrating significant HbA1c improvements and durable weight loss.^{31,32} Oral Semaglutide also demonstrated more flexible usage in weight control; however, it is not included in the NMA.³³ Liraglutide, though widely studied, is less potent than Semaglutide and is associated with higher rates of gastrointestinal adverse events, as shown in STEP-8 and corroborated in

observational cohorts.^{26,34} Cagrilintide, in contrast, is generally better tolerated but achieves smaller reductions in weight and metabolic outcomes, suggesting a niche role for patients prioritizing tolerability over maximum efficacy.³⁵

Safety outcomes across trials show broadly comparable rates of serious adverse events between active agents and placebo, consistent with prior GLP-1 receptor agonist meta-analyses.³⁶ However, gastrointestinal adverse events, particularly nausea, vomiting, and diarrhea, remain the primary tolerability issue and a major driver of discontinuation.³⁷ Higher discontinuation rates with semaglutide and liraglutide reflect the well-established dose-dependent relationship between incretin therapies and gastrointestinal side effects.³⁸ These symptoms often peak during dose escalation, especially at higher maintenance doses or in combination regimens.³⁹ Although structured titration schedules in trials help mitigate intolerance, real-world adherence to these protocols may be variable, contributing to greater symptom burden and higher discontinuation rates. Individuals such as older adults or those with multimorbidity may be especially vulnerable.

The findings of this analysis have important implications for clinical practice. Notably, most included trials enrolled relatively younger and healthier adults. The generalizability of these results to older adults, individuals with multimorbidity, and other clinically complex subgroups remains uncertain. These populations may exhibit different patterns of treatment response, tolerability, and adherence, highlighting the need for dedicated studies to better characterize the effectiveness and safety of these therapies in higher-risk groups. It is also important to situate these findings within the broader landscape of obesity pharmacotherapy. Several other agents, such as tirzepatide, phentermine/topiramate, naltrexone/bupropion, orlistat, and emerging incretin-based combinations, are widely used in clinical practice.^{40,41} Although comparisons with these therapies

were outside the scope of this NMA, clinicians should interpret the present results in the context of the full range of available treatment options. Treatment selection should be individualized, incorporating considerations of efficacy, tolerability, comorbidities, accessibility, and patient preference. Because long-term adherence is a key determinant of real-world effectiveness, proactive strategies to support persistence with therapy are essential for achieving and sustaining meaningful clinical outcomes.

Longer-term studies are needed to clarify the durability of weight loss, relapse rates, and maintenance strategies beyond two years. Cardiovascular and renal outcomes, which have been established for some GLP-1 RAs, must be evaluated specifically for dual agonists like CagriSema. Direct comparative studies with emerging agents, including tirzepatide and other multi-agonists, are also warranted to determine the optimal balance of efficacy and tolerability. Future research should incorporate real-world data to capture the effects of adherence, discontinuation, and cost, all of which may substantially affect effectiveness outside controlled trial conditions. Additionally, outcomes beyond weight and glycemia, such as quality of life, functional capacity, and patient-reported outcomes, should be more systematically integrated into trials to inform patient-centered care.

This analysis has several limitations. First, substantial heterogeneity across the included trials, encompassing differences in population characteristics, baseline comorbidities, and follow-up durations, may have influenced the validity of the indirect comparisons. Two studies with relatively short follow-up periods were also included; however, because they evaluated different agents, a robust sensitivity analysis could not be performed. Second, although no major inconsistency was identified, reporting of safety and treatment discontinuation outcomes varied

considerably among studies, limiting the precision of the pooled estimates. Third, many of the included trials were industry-sponsored, which may introduce reporting bias or selective emphasis on favorable outcomes. Finally, because this NMA is composed of indirect rather than direct comparisons, further head-to-head trials are required to strengthen the evidence base.

Just Accepted

359 **Conclusion**

360 This NMA confirms that CagriSema provides the greatest efficacy for body weight and
361 waist circumference reduction, with Semaglutide offering a strong balance of efficacy and
362 glycemic benefit. Cagrilintide demonstrates better tolerability with modest effects, while
363 Liraglutide is both less effective and less well tolerated. These findings underscore the central role
364 of incretin-based therapies in obesity management and highlight the need for individualized
365 treatment strategies. Future trials with extended follow-up and direct comparisons against
366 emerging therapies are necessary to further define optimal treatment approaches.

367

Acknowledgments: None.

Funding: None

Author Contributions: Y.M., and T.J. designed the study, conducted the literature search, performed the quality assessment, extracted data, carried out the analyses, and drafted the manuscript. L.R., M.L.T., and Y.W. contributed to data interpretation and critically revised the manuscript. All authors have read and approved the final version of the manuscript and agree with its content and data.

Data Availability Statement: The datasets used and analyzed in this study are available from the corresponding author upon reasonable request.

Ethics Statement: Institutional Review Board approval was waived, as this study is a meta-analysis of previously published data.

Conflicts of Interest: The authors declare no conflicts of interest.

381 Reference

- 382 1. World Health Organization. Obesity and overweight. [https://www.who.int/news-](https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight)
383 [room/fact-sheets/detail/obesity-and-overweight](https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight) (Published May 2025, Accessed on September 1,
384 2025)
- 385 2. Ahmed SK, Mohammed RA. Obesity: Prevalence, causes, consequences, management,
386 preventive strategies and future research directions. *Metabolism Open*. 2025/09/01/
387 2025;27:100375. doi:<https://doi.org/10.1016/j.metop.2025.100375>
- 388 3. Ullah MI, Tamanna S. Obesity: Clinical Impact, Pathophysiology, Complications, and
389 Modern Innovations in Therapeutic Strategies. *Medicines*. 2025;12(3):19.
- 390 4. Gostoli S, Raimondi G, Popa AP, Giovannini M, Benasi G, Rafanelli C. Behavioral
391 Lifestyle Interventions for Weight Loss in Overweight or Obese Patients with Type 2 Diabetes: A
392 Systematic Review of the Literature. *Curr Obes Rep*. Jun 2024;13(2):224-241.
393 doi:10.1007/s13679-024-00552-5
- 394 5. Pasarica M, Dhurandhar NV, Stensel DJ. Obesity management with next-generation drugs.
395 *Int J Obes (Lond)*. Mar 2025;49(3):367-368. doi:10.1038/s41366-025-01734-4
- 396 6. Moiz A, Fillion KB, Tsoukas MA, Yu OHY, Peters TM, Eisenberg MJ. The expanding role
397 of GLP-1 receptor agonists: a narrative review of current evidence and future directions.
398 *EClinicalMedicine*. Aug 2025;86:103363. doi:10.1016/j.eclim.2025.103363
- 399 7. Anam M, Maharjan S, Amjad Z, et al. Efficacy of Semaglutide in Treating Obesity: A
400 Systematic Review of Randomized Controlled Trials (RCTs). *Cureus*. Dec 2022;14(12):e32610.
401 doi:10.7759/cureus.32610
- 402 8. Amaro A, Skolnik NS, Sugimoto D. Cardiometabolic risk factors efficacy of semaglutide
403 in the STEP program. *Postgrad Med*. Jan 2022;134(sup1):18-27.
404 doi:10.1080/00325481.2022.2147325
- 405 9. Wharton S, Calanna S, Davies M, et al. Gastrointestinal tolerability of once-weekly
406 semaglutide 2.4 mg in adults with overweight or obesity, and the relationship between
407 gastrointestinal adverse events and weight loss. *Diabetes Obes Metab*. Jan 2022;24(1):94-105.
408 doi:10.1111/dom.14551
- 409 10. Lutz TA. Role of amylin in feeding and satiation. *Neuropharmacology*. 2025/11/01/
410 2025;278:110587. doi:<https://doi.org/10.1016/j.neuropharm.2025.110587>
- 411 11. Walker CS, Aitken JF, Vazhoor Amarsingh G, Zhang S, Cooper GJS. Amylin: emergent
412 therapeutic opportunities in overweight, obesity and diabetes mellitus. *Nat Rev Endocrinol*. Aug
413 2025;21(8):482-494. doi:10.1038/s41574-025-01125-9
- 414 12. D'Ascanio AM, Mullally JA, Frishman WH. Cagrilintide: A Long-Acting Amylin Analog
415 for the Treatment of Obesity. *Cardiology in Review*. 2024;32(1)
- 416 13. Reiss AB, Gulkarov S, Lau R, et al. v. *Biomolecules*. 2025;15(3):408.
- 417 14. Kokkorakis M, Chakhtoura M, Rhayem C, et al. Emerging pharmacotherapies for obesity:
418 A systematic review. *Pharmacological Reviews*. 2025/01/01/ 2025;77(1):100002.
419 doi:<https://doi.org/10.1124/pharmrev.123.001045>
- 420 15. Frias JP, Deenadayalan S, Erichsen L, et al. Efficacy and safety of co-administered once-
421 weekly cagrilintide 2·4 mg with once-weekly semaglutide 2·4 mg in type 2 diabetes: a multicentre,
422 randomised, double-blind, active-controlled, phase 2 trial. *Lancet*. Aug 26 2023;402(10403):720-
423 730. doi:10.1016/s0140-6736(23)01163-7

16. Bliddal H, Bays H, Czernichow S, et al. Once-Weekly Semaglutide in Persons with Obesity and Knee Osteoarthritis. *N Engl J Med*. Oct 31 2024;391(17):1573-1583. doi:10.1056/NEJMoa2403664
17. Enebo LB, Berthelsen KK, Kankam M, et al. Safety, tolerability, pharmacokinetics, and pharmacodynamics of concomitant administration of multiple doses of cagrilintide with semaglutide 2·4 mg for weight management: a randomised, controlled, phase 1b trial. *Lancet*. May 8 2021;397(10286):1736-1748. doi:10.1016/s0140-6736(21)00845-x
18. Garvey WT, Batterham RL, Bhatta M, et al. Two-year effects of semaglutide in adults with overweight or obesity: the STEP 5 trial. *Nat Med*. Oct 2022;28(10):2083-2091. doi:10.1038/s41591-022-02026-4
19. Garvey WT, Blüher M, Osorto Contreras CK, et al. Coadministered Cagrilintide and Semaglutide in Adults with Overweight or Obesity. *N Engl J Med*. Aug 14 2025;393(7):635-647. doi:10.1056/NEJMoa2502081
20. Davies MJ, Bajaj HS, Broholm C, et al. Cagrilintide-Semaglutide in Adults with Overweight or Obesity and Type 2 Diabetes. *N Engl J Med*. Aug 14 2025;393(7):648-659. doi:10.1056/NEJMoa2502082
21. Kadowaki T, Isendahl J, Khalid U, et al. Semaglutide once a week in adults with overweight or obesity, with or without type 2 diabetes in an east Asian population (STEP 6): a randomised, double-blind, double-dummy, placebo-controlled, phase 3a trial. *Lancet Diabetes Endocrinol*. Mar 2022;10(3):193-206. doi:10.1016/s2213-8587(22)00008-0
22. Lim S, Buranapin S, Bao X, et al. Once-weekly semaglutide 2·4 mg in an Asian population with obesity, defined as BMI $\geq$ 25 kg/m², in South Korea and Thailand (STEP 11): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Diabetes Endocrinol*. Aug 14 2025;doi:10.1016/s2213-8587(25)00164-0
23. Lincoff AM, Brown-Frandsen K, Colhoun HM, et al. Semaglutide and Cardiovascular Outcomes in Obesity without Diabetes. *N Engl J Med*. Dec 14 2023;389(24):2221-2232. doi:10.1056/NEJMoa2307563
24. McGowan BM, Bruun JM, Capehorn M, et al. Efficacy and safety of once-weekly semaglutide 2·4 mg versus placebo in people with obesity and prediabetes (STEP 10): a randomised, double-blind, placebo-controlled, multicentre phase 3 trial. *Lancet Diabetes Endocrinol*. Sep 2024;12(9):631-642. doi:10.1016/s2213-8587(24)00182-7
25. Mu Y, Bao X, Eliaschewitz FG, et al. Efficacy and safety of once weekly semaglutide 2·4 mg for weight management in a predominantly east Asian population with overweight or obesity (STEP 7): a double-blind, multicentre, randomised controlled trial. *Lancet Diabetes Endocrinol*. Mar 2024;12(3):184-195. doi:10.1016/s2213-8587(23)00388-1
26. Rubino DM, Greenway FL, Khalid U, et al. Effect of Weekly Subcutaneous Semaglutide vs Daily Liraglutide on Body Weight in Adults With Overweight or Obesity Without Diabetes: The STEP 8 Randomized Clinical Trial. *Jama*. Jan 11 2022;327(2):138-150. doi:10.1001/jama.2021.23619
27. Wilding JPH, Batterham RL, Calanna S, et al. Once-Weekly Semaglutide in Adults with Overweight or Obesity. *N Engl J Med*. Mar 18 2021;384(11):989-1002. doi:10.1056/NEJMoa2032183
28. Ryan DH, Yockey SR. Weight Loss and Improvement in Comorbidity: Differences at 5%, 10%, 15%, and Over. *Curr Obes Rep*. Jun 2017;6(2):187-194. doi:10.1007/s13679-017-0262-y

29. Bhat S, Fernandez CJ, Lakshmi V, Pappachan JM. Efficacy and safety of incretin co-agonists: Transformative advances in cardiometabolic healthcare. *World J Cardiol.* Aug 26 2025;17(8):107991. doi:10.4330/wjc.v17.i8.107991
30. Krüger N, Schneeweiss S, Fuse K, et al. Semaglutide and Tirzepatide in Patients With Heart Failure With Preserved Ejection Fraction. *Jama.* Aug 31 2025;doi:10.1001/jama.2025.14092
31. Thomsen RW, Mailhac A, Løhde JB, Pottegård A. Real-world evidence on the utilization, clinical and comparative effectiveness, and adverse effects of newer GLP-1RA-based weight-loss therapies. *Diabetes Obes Metab.* Apr 2025;27 Suppl 2(Suppl 2):66-88. doi:10.1111/dom.16364
32. Moiz A, Filion KB, Toutounchi H, et al. Efficacy and Safety of Glucagon-Like Peptide-1 Receptor Agonists for Weight Loss Among Adults Without Diabetes : A Systematic Review of Randomized Controlled Trials. *Ann Intern Med.* Feb 2025;178(2):199-217. doi:10.7326/annals-24-01590
33. Wharton S, Lingvay I, Bogdanski P, et al. Oral Semaglutide at a Dose of 25 mg in Adults with Overweight or Obesity. *N Engl J Med.* Sep 18 2025;393(11):1077-1087. doi:10.1056/NEJMoa2500969
34. Stefanou MI, Chatziralli I, Lambadiari V, et al. Efficacy and Safety of GLP-1 and Dual GIP/GLP-1 Receptor Agonists in Idiopathic Intracranial Hypertension: A Systematic Review and Meta-Analysis. *Eur J Neurol.* Sep 2025;32(9):e70358. doi:10.1111/ene.70358
35. Carvas AO, Leuthardt A, Kulka P, et al. Cagrilintide lowers bodyweight through brain amylin receptors 1 and 3. *EBioMedicine.* Aug 2025;118:105836. doi:10.1016/j.ebiom.2025.105836
36. Njei B, Al-Ajlouni Y, Lemos SY, et al. Efficacy and Safety of GLP-1 Receptor Agonists in Patients With Metabolic Dysfunction-Associated Steatotic Liver Disease: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Cureus.* Oct 2024;16(10):e71366. doi:10.7759/cureus.71366
37. Ismaiel A, Scarlata GGM, Boitos I, et al. Gastrointestinal adverse events associated with GLP-1 RA in non-diabetic patients with overweight or obesity: a systematic review and network meta-analysis. *International Journal of Obesity.* 2025/10/01 2025;49(10):1946-1957. doi:10.1038/s41366-025-01859-6
38. Galli M, Benenati S, Laudani C, et al. Cardiovascular Effects and Tolerability of GLP-1 Receptor Agonists: A Systematic Review and Meta-Analysis of 99,592 patients. *J Am Coll Cardiol.* Aug 19 2025;doi:10.1016/j.jacc.2025.08.027
39. Aroda VR, Bain SC, Cariou B, et al. Efficacy and safety of once-weekly semaglutide versus once-daily insulin glargine as add-on to metformin (with or without sulfonylureas) in insulin-naïve patients with type 2 diabetes (SUSTAIN 4): a randomised, open-label, parallel-group, multicentre, multinational, phase 3a trial. *The Lancet Diabetes & Endocrinology.* 2017;5(5):355-366. doi:10.1016/S2213-8587(17)30085-2
40. Liu L, Li Z, Ye W, et al. Safety and effects of anti-obesity medications on weight loss, cardiometabolic, and psychological outcomes in people living with overweight or obesity: a systematic review and meta-analysis. *eClinicalMedicine.* 2025;79doi:10.1016/j.eclinm.2024.103020
41. Gudzone KA, Kushner RF. Medications for Obesity: A Review. *Jama.* Aug 20 2024;332(7):571-584. doi:10.1001/jama.2024.10816

Table 1. Characteristics and Background of Enrolled Studies

Author	Inclusion criteria	Treatment arms	Cases	Age (year)	Female (%)	BW (kg)	BMI (kg/m ²)	CW (cm)	HbA1c (%)	Follow up
Bliddal et al., 2024	BMI > 30 with osteoarthritis	Semaglutide vs. Placebo	407	56 (10)	332 (82)	109 (24)	40.3 (7.2)	119 (16)	–	68w
Enebo et al., 2021	BMI 27.0–39.9 kg/m ²	CagriSema vs. Semaglutide	36	42 (9)	15 (42)	97 (15)	32.2 (2.8)	–	5.3 (0.4)	20w
Garvey et al., 2022	BMI ≥ 27 with complication or BMI ≥ 30	Semaglutide vs. Placebo	304	47 (11)	236 (78)	106 (22)	38.6 (6.9)	116 (15)	5.7 (0.4)	104w
Garvey et al., 2025	BMI ≥ 27 with complication or BMI ≥ 30	CagriSema vs. Semaglutide vs. Cagrilintide vs Placebo	3,417	47 (12)	2309 (68)	107 (23)	37.9 (6.8)	115 (16)	5.5 (0.4)	68w
Kadowaki et al., 2022 ^a	BMI ≥ 27 with complication or BMI ≥ 35	Semaglutide vs. Placebo	300	51 (11)	111 (37)	88 (16)	32 (4.5)	104 (11)	6.4 (1.2)	68w
Lau et al., 2021	BMI ≥ 27 with complication or BMI ≥ 30	Cagrilintide vs. Placebo vs. Liraglutide	302	52 (10)	199 (66)	107 (23)	37.9 (6.9)	114 (14)	5.6 (0.4)	26w
Lim et al., 2025	BMI ≥ 25	Semaglutide vs. Placebo	150	39 (11)	111 (74)	84 (18)	31.3 (5.2)	98 (12)	5.6 (0.3)	44w
Lincoff et al., 2023	BMI ≥ 27 with preexisting cardiovascular disease	Semaglutide vs. Placebo	17,604	62 (9)	4892 (28)	97 (18)	33.3 (5.0)	111 (13)	5.8 (0.3)	48w
McGowan et al., 2024	BMI > 30 with prediabetes	Semaglutide vs. Placebo	207	53 (11)	147 (71)	112 (22)	40.1 (6.9)	120 (15)	5.9 (0.3)	52w
Mu et al., 2024 ^b	BMI ≥ 27 with complication or BMI ≥ 30	Semaglutide vs. Placebo	375	40 (11)	170 (45)	96 (18)	34 (4.8)	108 (11)	6.3 (1.1)	44w
Rubino et al., 2022	BMI ≥ 27 with complication or BMI ≥ 31	Semaglutide vs. Liraglutide vs. Placebo	338	49 (13)	265 (78)	105 (24)	37.5 (6.8)	113 (16)	5.5 (0.3)	68w
Wilding et al., 2021	BMI ≥ 27 with complication or BMI ≥ 30	Semaglutide vs. Placebo	1,961	46 (13)	1453 (74)	105 (22)	37.9 (6.6)	115 (15)	5.7 (0.3)	68w

a. 99 (25%) patients were diagnosed with type 2 diabetes mellitus; b. 96 (25.6%) patients were diagnosed with type 2 diabetes mellitus

BW: body weight; BMI: body mass index; WC: waist circumference; w: weeks; vs: versus

Table 2. Network Meta-Analysis of Body Weight and Waist Circumference Changes

Body Weight [#]					
CagriSema					
6.6 (3.0, 10.3)	Semaglutide				
11.8 (6.5, 17.5)	5.2 (1.0, 9.7)	Cagrilintide			
13.4 (8.8, 18.5)	6.8 (3.5, 10.4)	1.6 (-2.6, 5.8)	Liraglutide		
17.7 (14.2, 21.3)	11.1 (9.5, 12.7)	5.9 (1.5, 10.0)	4.3 (0.7, 7.6)	Placebo	
Waist Circumference					
CagriSema					
4.9 (0.9, 8.9)	Semaglutide				
9.7 (4.8, 15.0)	4.8 (1.4, 8.7)	Cagrilintide			
11.1 (6.7, 16.0)	6.2 (3.5, 9.4)	1.4 (-1.9, 5.0)	Liraglutide		
13.4 (9.7, 17.1)	8.4 (7.0, 10.0)	3.6 (0.0, 7.0)	2.3 (-0.9, 4.9)	Placebo	
Serious Adverse Events [#]					
Semaglutide					
-0.4 (-1.0, 0.3)	Cagrilintide				
-0.5 (-1.2, 0.3)	-0.1 (-1.0, 0.9)	Liraglutide			
-0.5 (-1.1, 0.1)	-0.1 (-0.8, 0.6)	-0.1 (-1.0, 0.9)	CagriSema		
-0.0 (-0.3, 0.3)	0.4 (-0.3, 1.0)	0.5 (-0.3, 1.2)	0.5 (-0.1, 1.1)	Placebo	
Treatment Discontinuation					
CagriSema					
0.1 (-1.4, 1.9)	Liraglutide				
0.2 (-1.0, 1.6)	0.1 (-1.1, 1.3)	Semaglutide			
0.7 (-0.7, 2.1)	0.6 (-0.9, 1.9)	0.4 (-0.7, 1.6)	Cagrilintide		
0.9 (-0.3, 2.2)	0.8 (-0.4, 1.9)	0.6 (0.1, 1.2)	0.2 (-0.9, 1.3)	Placebo	
Treatment Discontinuation Due to Gastrointestinal Symptoms					
Liraglutide					
0.9 (-1.3, 3.3)	CagriSema				
1.7 (-0.2, 3.9)	0.8 (-0.2, 2.0)	Semaglutide			
2.1 (-0.3, 4.8)	1.1 (-0.2, 2.8)	0.3 (-1.1, 2.0)	Cagrilintide		
3.3 (1.5, 5.6)	2.5 (1.3, 3.7)	1.7 (1.0, 2.3)	1.3 (-0.4, 2.9)	Placebo	


[#]: Cells highlighted in yellow indicate statistically significant differences.

Table 3. Summary of Certainty of Evidence

Outcomes	Impact (Relative to Placebo)	No. of Participants (Studies)	Certainty of the Evidence (GRADE)
Body weight reduction	CagriSema (MD: -17.7%), Semaglutide (MD: -11.9%), Cagrilintide (MD: -5.9%), Liraglutide (MD: -4.3%)	25,401 (12 studies)	⊕⊕⊕⊕ High
Waist circumference reduction	CagriSema (MD: -13.4 cm), Semaglutide (MD: -8.4 cm), Cagrilintide (MD: -3.6 cm), Liraglutide (MD: -2.3 cm)	24,958 (10 studies)	⊕⊕⊕⊕ High
HbA1c reduction	Semaglutide (MD: -0.45%)	18,923 (5 studies)	⊕⊕⊕⊕ High
Serious adverse events	No significant increase observed (OR range: 1.0–1.7)	25,401 (12 studies)	⊕⊕⊕⊖ Moderate
Treatment discontinuation	Semaglutide was associated with an increased risk (OR: 1.9), while other agents showed no statistically significant differences.	25,401 (12 studies)	⊕⊕⊕⊖ Moderate
Treatment discontinuation due to gastrointestinal symptoms	Cagrilintide showed no statistically significant difference	24742 (9 studies)	⊕⊕⊕⊖ Moderate


No.: Number; GRADE: Grading of Recommendations, Assessment, Development, and Evaluation; MD: Mean Difference; OR: Odds Ratio

Figure 1. Network graph of enrolled studies



A: Body weight; B: Waist circumference; C: Serious adverse events and treatment discontinuation; D: Treatment discontinuation due to gastrointestinal symptoms

Figure 2. Effects of each agent compared with placebo across different outcomes



A: Body weight; B: Waist circumference; C: Serious adverse events; D: Treatment discontinuation; E: Discontinuation due to gastrointestinal symptoms; F: HbA1c reduction.