

Type of Article: Original Research

Title: Efficacy of Antihypertensive Therapy in Primary IgA Nephropathy in Adults: A Systematic Review and Network Meta-Analysis of Randomized Controlled Trials

Authors: Yoshihito Iida¹, Guihua Gao², Chadia Beaini³, Ali Ghanem³, Sandeep Varma⁴, Magdy Elsharkawy⁵

¹ Department of Nephrology, Okanishi Clinic, Shimada, Japan

² Department of Nephrology, Oeda Internal Medicine Clinic, Toyohashi, Japan

³ Department of Nephrology, Mediclinic Middle East, Dubai, United Arab Emirates

⁴ Department of Nephrology, Prime Hospital, Dubai, United Arab Emirates

⁵ Department of Nephrology, Ain-Shams University, Cairo, Egypt

Correspondence: Magdy Elsharkawy

Address: 1 Elsarayat St., Abbaseya, Cairo, Egypt, 11517

Department of Nephrology, Ain-Shams University

Email: magdi35@hotmail.com

Tel: (+20) 26831231

ORCID: <https://orcid.org/0000-0002-2007-7385>

19 Is antihypertensive treatment recommended for patients with primary IgA nephropathy in adults?

20 Moderate-certainty evidence supports antihypertensive agents, particularly angiotensin-
21 converting enzyme inhibitors, angiotensin II receptor blockers, and endothelin receptor antagonists,
22 for reducing proteinuria in primary IgA nephropathy. However, there is only low-certainty
23 evidence regarding their effectiveness in improving the estimated glomerular filtration rate and
24 serum creatinine levels. Additionally, strong certainty of evidence indicates an increased risk of
25 adverse events associated with these treatments. Nevertheless, antihypertensive therapy is strongly
26 recommended for managing IgA nephropathy due to its proven benefits in reducing proteinuria
27 and slowing disease progression.

Background: IgA nephropathy (IgAN) is a leading cause of chronic kidney disease, with proteinuria and hypertension accelerating the disease's progression. While the renin-angiotensin-aldosterone system inhibitors remain the standard treatment, emerging therapies such as endothelin receptor antagonists (ERAs) offer potential renoprotective benefits. This study conducted a network meta-analysis (NMA) of randomized controlled trials (RCTs) to compare the efficacy of various antihypertensive agents in reducing proteinuria and preserving kidney function in adults with IgAN.

Methods: A comprehensive literature search was performed on the PubMed, Cochrane Library, and Web of Science databases through February 28, 2025. RCTs evaluating angiotensin-converting enzyme inhibitors (ACEIs), angiotensin II receptor blockers (ARBs), ERAs, calcium channel blockers, and direct renin inhibitors in adults with biopsy-confirmed IgAN were reviewed in the study. A frequentist random-effects NMA was conducted to assess treatment efficacy in reducing proteinuria, improving estimated glomerular filtration rate (eGFR), and lowering serum creatinine (Scr) levels.

Results: Thirteen RCTs, which covered 1,572 patients, were analyzed. Sparsentan exhibited the most significant reduction in proteinuria (mean difference [MD]: 2.42; 95% confidence interval [CI]: 0.06, 4.79), followed by Benazepril (MD: 1.92; 95% CI: 0.32, 3.52) and Irbesartan (MD: 1.63; 95% CI: -0.51, 3.78). However, not all studies reported outcomes regarding eGFR and Scr. In studies reporting eGFR outcomes, Sparsentan ranked highest for improvement (MD: 12.6; 95% CI: -30.7, 55.8), followed by Irbesartan (MD: 8.2; 95% CI: -29.8, 46.1) and Enalapril (MD: 6.2; 95% CI: -21.4, 33.7). Enalapril demonstrated the most significant Scr reduction (MD: 0.9; 95%

CI: -0.07, 1.73), followed by Benazepril (MD: 0.74; 95% CI: 0.32, 1.16) and Losartan (MD: 0.18; 95% CI: -0.11, 0.47).

Conclusion: This NMA confirms that ACEIs and ARBs are suitable first-line treatments for IgAN while highlighting ERAs, particularly Sparsentan, as a promising alternative for patients with persistent proteinuria.

Keywords: IgA nephropathy, antihypertensive therapy, proteinuria, endothelin receptor antagonists, network meta-analysis.

Introduction

IgA nephropathy (IgAN) is the leading cause of primary glomerulonephritis. Its prevalence is highest among East Asian and White populations, though it remains relatively uncommon in Black individuals.¹ IgAN can be diagnosed only upon evaluation of a kidney biopsy with immunofluorescence microscopy. Globally, the incidence of IgAN varies from 0.06 in South Africa to 4.2 per 100,000 in Japan.² A meta-analysis of US studies calculated an annual IgAN Incidence of 1.29 per 100,000 people, translating to a yearly incidence of the disease in 4236 adults and children in the US.³ IgAN is a significant cause of chronic kidney disease (CKD) and can lead to a progressive decline in kidney function. The severity and progression of IgAN are influenced by proteinuria, hypertension, and histological findings.⁴ A study analyzing data from the United States Renal Data System reported that patients with IgAN-attributed kidney failure incur annual healthcare costs exceeding \$63,000 per patient.⁵ Patients with high-risk proteinuria (≥ 1 g/day) and worsening kidney function incur higher healthcare costs, with expenses rising as CKD progresses. Their monthly costs average \$3,732, compared to \$1,457 for those with lower proteinuria. Moreover, CKD-related costs jump from \$2,111 in Stage 1 of the disease to \$10,703 in Stage 5.⁶

Over the past few decades, therapeutic approaches for IgAN have evolved significantly, transitioning from conventional supportive care to more targeted interventions to reduce proteinuria and slow the disease's progression.⁷ Nevertheless, the cornerstone of the therapy remains the blockade of the renin-angiotensin-aldosterone System (RAAS), primarily through angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin II receptor blockers (ARBs), which have consistently demonstrated reduced proteinuria and preserved renal function.⁸ More recently, novel therapeutic agents have expanded the treatment landscape for IgAN, offering

additional renoprotective benefits. Sodium-glucose cotransporter-2 (SGLT2) inhibitors, initially developed for diabetes management, have shown promise in reducing proteinuria and mitigating the decline in estimated glomerular filtration rate (eGFR) in IgAN patients, even those without diabetes.⁹ Endothelin receptor antagonists (ERAs), targeting the endothelin pathway involved in renal fibrosis and inflammation, have emerged as another promising approach for high-risk patients.¹⁰ Additionally, corticosteroid-sparing immunosuppressive agents are being investigated as alternatives to traditional immunosuppressive regimens, potentially reducing the adverse effects of long-term steroid use.¹¹

Hypertension is a key modifiable risk factor in IgAN and is critical to the disease's progression. Elevated blood pressure exacerbates glomerular hypertension, accelerating tubulointerstitial fibrosis and renal function decline.⁴ Clinical trials have shown that effective blood pressure control significantly reduces proteinuria and slows the decline of eGFR in IgAN patients.¹² While RAAS blockade remains the first-line therapy for IgAN, studies on the disease have explored additional antihypertensive agents, such as calcium channel blockers (CCBs) and direct renin inhibitors, for their potential benefits.¹³ Despite various antihypertensive strategies, the optimal treatment approach remains uncertain, underscoring the need for a comprehensive comparative analysis.

However, there are still few direct head-to-head comparisons between these emerging therapies, and the most effective treatment strategies for specific patient subgroups remain unclear. Emerging evidence suggests that ERAs provide additional renoprotective benefits beyond conventional antihypertensive therapies, highlighting their potential role in IgAN management.¹⁴ Given the heterogeneity of treatment options and patient responses, a network meta-analysis

(NMA) is warranted to compare the efficacy and safety of different antihypertensive regimens systematically. Unlike traditional pairwise meta-analyses, NMAs integrate data from multiple randomized controlled trials (RCTs) to establish a hierarchical ranking of the analyzed therapies. By incorporating ERAs into this framework, the current NMA aims to optimize individualized treatment strategies and improve clinical decision-making in IgAN.

Just Accepted

109 **Methods**

110 *Protocol and Registration*

111 This NMA was registered in the Open Science Framework.¹⁵ The study protocol adhered
112 to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020
113 guidelines and was reviewed before analysis to ensure methodological rigor and transparency.¹⁶

114 *Eligibility Criteria*

115 The inclusion criteria for this NMA were as follows: 1) adult patients with biopsy-
116 confirmed IgAN; 2) studies focusing on patients with proteinuria ≥ 0.5 g/24h; 3) investigations of
117 interventions such as RAAS blockers, CCBs, direct renin inhibitors (DRIs), ERAs, or placebo; and
118 4) inclusion of patients regardless of hypertension. Exclusion criteria included the following: 1)
119 studies involving add-on therapy or immunosuppressive therapy, 2) phase 2 trials, 3) studies
120 incorporating tonsillectomy, 4) studies with insufficient data, and 5) observational study designs.
121 To minimize selection bias, no language restrictions were applied.

122 *Information Sources and Search Strategy*

123 A comprehensive literature search was conducted on the PubMed, the Cochrane Library,
124 and the Web of Science databases from inception to February 28, 2025. The search strategy
125 incorporated controlled vocabulary and relevant keywords to identify targeted patients, including
126 “IgA nephropathy”; interventions, including “hypertension,” “angiotensin-converting enzyme
127 inhibitors,” “angiotensin receptor blockers,” “calcium channel blockers,” “direct renin inhibitors,”

and “endothelin receptor antagonists”; and study design, including “randomized” and “controlled.”
Additionally, the reference lists of relevant reviews and included studies were screened manually.

Selection Process

Two independent reviewers (Y.I. and G.G.) screened the titles and abstracts of the retrieved records. Full-text articles of potentially eligible studies were assessed per the inclusion criteria. Disagreements were resolved through discussion or consultation with a third reviewer (M.E.). The selection process was documented using a PRISMA flow diagram, which detailed the number of included and excluded studies at each stage and the reasons for exclusion (Figure S1).

Data Collection Process

A standardized data extraction form collected key study details, including the first author, publication year, country, sample size, and follow-up duration. Patient characteristics, such as age, sex, baseline serum creatinine (Scr), eGFR, and inclusion criteria, were recorded. Intervention details included the type of drug, dosage, and treatment duration. Outcomes assessed comprised changes in proteinuria, eGFR, and Scr levels, and reported adverse events (AEs). A second reviewer independently verified all extracted data to ensure accuracy. Several studies identified usual care primarily through the use of CCBs, and in the data analysis, CCBs were combined with usual care into a single category.

Outcomes

The primary outcomes varied across the selected studies; however, all reported proteinuria. The primary outcome was reduced proteinuria, which was measured as the mean difference from

baseline. Secondary outcomes included the decline in eGFR, changes in Scr levels, and the incidence of AEs.

Statistical Analysis

A frequentist random-effects NMA was conducted using the “meta” and “netmeta” packages in R. Results were reported as mean differences (MDs) or odds ratios (ORs) with 95% confidence intervals (CIs), and a two-tailed p-value < 0.05 was considered statistically significant. Treatment efficacy was ranked with the surface under the cumulative ranking curve (SUCRA). Heterogeneity was assessed using the I^2 statistic and Cochran’s Q test, while inconsistency was evaluated through node-splitting analyses. The sensitivity analyses excluded high-risk-of-bias studies. If substantial heterogeneity ($I^2 > 50\%$) was observed, subgroup analyses would be conducted based on baseline proteinuria levels or hypertension status. The publication bias was assessed using funnel plots and Egger’s tests.

Risk of Bias and Certainty of Evidence

The methodological quality of the included RCTs was assessed using the Cochrane Risk of Bias 2 tool, which evaluates aspects such as randomization, blinding, incomplete outcome data, and selective reporting.¹⁷ Two independent reviewers conducted the assessment, while a third reviewer resolved the discrepancies. The certainty of evidence was evaluated using the Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) approach, classifying it as high, moderate, low, or very low based on factors such as risk of bias, inconsistency, indirectness, imprecision, and publication bias.¹⁸

Results

Background of Studies

Eight hundred sixty-six articles were identified through database searches for the study's purpose. After 115 duplicate articles were removed, 662 more were excluded in the first screening and 76 in the second (Figure S1). Ultimately, 13 RCTs were included in the final analysis, comprising 1,572 participants with IgAN (Table 1).¹⁹⁻³¹ These studies were conducted across multiple countries, including Australia, Hong Kong, Italy, Japan, Korea, China, Spain, Singapore, and international multicenter trials. Their sample sizes ranged from 20 to 404 participants, whose mean age varied across studies, ranging from 29 to 51 years. The proportion of male participants was relatively high, ranging from 20% to 74%. Scr levels, when reported, varied widely from 0.8 mg/dL to 1.5 mg/dL, while eGFR values ranged from 58 mL/min/1.73m² to 111 mL/min/1.73m².

This NMA identified three specific CCBs (Amlodipine, Nifedipine, Verapamil), four types of ACEIs (Benazepril, Enalapril, Fosinopril, and Trandolapril), four kinds of ARBs (Candesartan, Irbesartan, Losartan, and Valsartan), two types of ERAs (Atrasentan and Sparsentan), and one DRI (Aliskiren). CCB studies were merged into the usual care group, as three previous studies had primarily defined usual care as CCB use. Two studies included patients without hypertension, three included those with hypertension, and others did not specify hypertension status. The use of add-on antihypertensive therapy varied across studies; five studies did not include add-on antihypertensives, while one study combined ACEI or ARB in all patients. Follow-up periods varied widely, from 1 month to 6 years.

188 *Proteinuria Reduction*

189 The network graph of the enrolled studies assessing proteinuria is presented in Figure S2.
190 Six studies used a placebo as the control treatment. Each agent is directly compared with the
191 placebo in Figure 2A. Sparsentan demonstrated the greatest reduced proteinuria, with a MD of
192 2.42 (95% CI: 0.06, 4.79), followed by Benazepril (MD: 1.92, 95% CI: 0.32, 3.52), Irbesartan
193 (MD: 1.63, 95% CI: -0.51, 3.78), Enalapril (MD: 1.41, 95% CI: 0, 2.83), Losartan (MD: 1.33, 95%
194 CI: -0.03, 2.69), Trandolapril (MD: 0.80, 95% CI: -0.37, 1.97), Candesartan (MD: 0.80, 95% CI:
195 -0.40, 2.00), Atrasentan (MD: 0.69, 95% CI: -0.32, 1.70), usual care (MD: 0.50, 95% CI: -0.66,
196 1.66), Aliskiren (MD: 0.50, 95% CI: -0.77, 1.77), Fosinopril (MD: 0.42, 95% CI: -0.75, 1.59), and
197 Valsartan (MD: 0.19, 95% CI: -0.94, 1.32).

198 The NMA results for proteinuria reduction are presented in Table 2. Although Sparsentan
199 showed the most significant decrease in proteinuria in direct comparisons, its superiority over
200 Benazepril, Irbesartan, Enalapril, Losartan, Trandolapril, Candesartan, Atrasentan, Aliskiren,
201 Fosinopril, usual care, and Valsartan was not statistically significant. The rank of efficiency based
202 on 1,000 simulations is depicted in Figure S3, with Sparsentan (0.880), Benazepril (0.827),
203 Irbesartan (0.723), Enalapril (0.710), Losartan (0.684), Candesartan (0.464), Trandolapril (0.459),
204 Atrasentan (0.395), Aliskiren (0.343), Usual care (0.326), Fosinopril (0.323), Valsartan (0.235),
205 and placebo (0.132). Cochran's Q statistic was used to assess heterogeneity and inconsistency, with
206 a total Q value of 9.310 (df = 3, p = 0.025), within-design Q value of 3.340 (df = 2, p = 0.188),
207 and between-design Q value of 5.97
208 0 (df = 1, p = 0.015).

209 *eGFR Improvement*

210 The network graph of the enrolled studies assessing proteinuria is presented in Figure S4.
211 Seven studies used a placebo as the control. Three studies used a placebo as the control treatment,
212 including one evaluating the ERA Atrasentan. The direct comparison of each agent with placebo
213 is presented in Figure 1B. Sparsentan demonstrated the most significant reduction in proteinuria,
214 with an MD of 12.6 (95% CI: -30.7, 55.8), followed by Irbesartan (MD: 8.2, 95% CI: -29.8, 46.1),
215 Enalapril (MD: 6.2, 95% CI: -21.4, 33.7), and Losartan (MD: 3.73, 95% CI: -24.1, 31.5).

216 The NMA results for proteinuria reduction are presented in Table 3. Although Sparsentan
217 showed the most significant improvement in eGFR in direct comparisons, its superiority over
218 Irbesartan, Enalapril, Trandolapril, Losartan, and Valsartan was not statistically significant. The
219 rank of efficiency in Scr reduction based on SUCRA values is shown in Figure S5, with Enalapril
220 (0.923), Benazepril (0.892), Losartan (0.604), usual care (0.477), placebo (0.263), Trandolapril
221 (0.254), and Candesartan (0.088). The Q statistics to assess homogeneity and consistency showed
222 a total Q of 0.18 (df = 1, p = 0.671), with within-design heterogeneity at Q = 0.18 (df = 1, p =
223 0.671) and no between-design heterogeneity (Q = 0.00, df = 0).

224 *Reducing Scr*

225 The network graph of the enrolled studies assessing Scr is presented in Figure S6. Three
226 studies used a placebo as the control treatment. Each agent is directly compared with the placebo
227 in Figure 1C. Enalapril demonstrated the most significant reduction in Scr, with an MD of 0.9 (95%
228 CI: -0.07, 1.73), followed by Benazepril (MD: 0.74; 95% CI: 0.32, 1.16), Losartan (MD: 0.18; 95%
229 CI: -0.11, 0.47), and usual care (MD: 0.10; 95% CI: -0.13, 0.33).

The NMA results for Scr reduction are presented in Table 4. Enalapril showed statistical superiority over Trandolapril and Candesartan with reductions of MDs of 0.90 (95% CI: 0.07, 1.73) and 1.00 (95% CI: 0.18, 1.82), respectively. The rank of efficiency in eGFR based on SUCRA values is shown in Figure S5, with Sparsentan (0.686), Irbesartan (0.643), Enalapril (0.610), Trandolapril (0.562), Losartan (0.545), Valsartan (0.465), Candesartan (0.448), placebo (0.445), Fosinopril (0.433), usual care (0.360), and Aliskiren (0.303). The Q statistics to assess homogeneity and consistency showed a total Q of 14.26 (df = 3, p = 0.003), with within-design heterogeneity at Q = 6.30 (df = 2, p = 0.043) and between-design heterogeneity at Q = 7.96 (df = 1, p = 0.005).

AEs

Due to the limited number of studies, NMA could not be conducted for AE assessment. Instead, a meta-analysis evaluated Atrasentan, Valsartan, and Aliskiren compared to placebo, yielding an odds ratio of 0.93 (95% CI: 0.58–1.48, p = 0.122; I² = 52.3%, Figure S8).

Bias and Certainty of Evidence

Egger's test for proteinuria reduction, eGFR, Scr reduction, and AEs yielded p-values of 0.388, 0.062, 0.142, and 0.473, respectively, indicating no significant evidence of publication bias. Heat plots for proteinuria reduction and Scr were presented in Figures S9 and S10, respectively. Notably, the comparison between Enalapril and Losartan demonstrated a higher degree of inconsistency. The risk of bias assessment is presented in Figure S11, showing that two studies had a high risk, seven had some concerns, and four had a low risk of bias. According to the GRADE approach, the certainty of evidence for proteinuria reduction was rated as moderate and downgraded due to the risk of bias. The certainty of evidence for eGFR was rated as low,

252 downgraded due to heterogeneity and risk of bias. At the same time, the certainty of evidence for
253 Scr was also rated as low, downgraded due to the risk of bias and imprecision.

Just Accepted

Discussion

This NMA comprehensively evaluated the efficacy of various antihypertensive agents in IgAN, providing valuable insights into their relative renoprotective effects. Among these agents, Sparsentan emerged as the most effective in reducing proteinuria and preserving eGFR; however, its superiority over other treatments could not be statistically confirmed. Benazepril, Enalapril, and Irbesartan ranked highest among ACEIs and ARBs for proteinuria reduction and eGFR improvement, reinforcing their well-established role in IgAN management. Additionally, other ACEIs and ARBs demonstrated greater renoprotective effects than usual care, which primarily involved CCBs, further supporting their preferential use in clinical practice. In contrast, DRIs exhibited relatively lower effectiveness in renoprotection, suggesting that they play a more limited role in IgAN treatment. Within ERAs, differences in efficacy were observed between Sparsentan and Atrasentan, underscoring the need for further investigation into class-specific variations. Notably, no significant safety concerns were identified, although the available data on AEs were limited. These findings are consistent with previous studies emphasizing the benefits of ACEIs and ARBs while highlighting ERAs, particularly Sparsentan, as a promising emerging therapeutic option for IgAN.³²

The findings of this NMA have important clinical implications for IgAN management. The consistent renoprotective benefits observed with ACEIs and ARBs reinforce their role as first-line therapy, in line with current guidelines.^{33,34} Benazepril, Enalapril, and Irbesartan, ranked highest within their respective classes, may be preferred for maximizing proteinuria reduction and eGFR preservation. The superior efficacy of Sparsentan, despite the lack of statistically confirmed superiority over other agents, highlights its potential as an emerging treatment option, particularly given its dual inhibition of the endothelin and angiotensin pathways. **The absence of statistical**

significance may be attributed to limited sample sizes, overlapping confidence intervals, and variations in study design and patient characteristics across included trials.

The observed differences in efficacy among ERAs suggest further research to establish their optimal use in IgAN treatment. Additionally, the relatively lower effectiveness of DRIs suggests that they may play a limited role in clinical practice. Importantly, no major safety concerns were identified, supporting these treatments' feasibility in real-world settings. These findings align with previous research demonstrating the renoprotective effects of RAAS blockade while introducing ERAs as a promising adjunctive strategy for patients at high risk of progression despite standard therapies.

The therapeutic effects of antihypertensive agents in IgAN extend beyond blood pressure control; they target key pathophysiological mechanisms involved in disease progression.³⁵ ACEIs and ARBs remain the cornerstone of treatment by inhibiting the RAAS, reducing intraglomerular pressure, and exerting anti-inflammatory and antifibrotic effects.³⁶ Their superior ranking in this NMA reinforces their role as first-line therapy treatments. ERAs, such as Sparsentan and Atrasentan, target the endothelin system, with Sparsentan offering dual RAS and endothelin-1 inhibition, potentially enhancing renoprotection. Differences in efficacy between ERAs warrant further investigation. DRIs, which act upstream in the RAS, demonstrate relatively lower efficacy, possibly due to the compensatory mechanisms limiting their renoprotective impact.³⁷ While CCBs aid blood pressure control, their direct renoprotective effects appear inferior to RAS inhibitors.

Combination therapy in IgAN presents a promising approach to enhancing renoprotection by targeting multiple pathophysiological mechanisms. Combining ERAs, particularly Sparsentan, with ACEIs or ARBs may reduce proteinuria by addressing the angiotensin II and endothelin-1 pathways.³⁸ Similarly, SGLT2 inhibitors have emerged as adjunctive therapies that reduce

glomerular hyperfiltration and inflammation, demonstrating significant renoprotective effects in combination with RAAS inhibitors.³⁹ For high-risk IgAN patients, corticosteroids or immunosuppressants may be used alongside RAAS inhibition, as evidenced by the TESTING trial. However, careful patient selection is necessary due to potential adverse effects.⁴⁰ Combination strategies should be tailored to disease severity and individual patient risk, and further studies are needed to refine their long-term efficacy and safety.

This study has several limitations that need to be considered when interpreting the findings. First, although NMA allows indirect comparisons between treatments, the inherent variability in study designs, patient populations, and treatment regimens may introduce heterogeneity, which can affect the robustness of the results. **Second, while this analysis identified Sparsentan as the most effective proteinuria reduction and eGFR preservation agent, its superiority over other treatments was not statistically confirmed. This highlights the need for more direct head-to-head trials.** Third, data on combination therapies, particularly ERAs with RAAS or SGLT2 inhibitors, remain limited, necessitating further clinical studies to establish their optimal use and long-term effects. Additionally, due to the limited availability of safety data, especially for newer agents such as Sparsentan and Atrasentan, a comprehensive assessment of the risk-benefit profiles of these agents cannot be conducted. **Finally, due to the limited availability of safety data across included trials, a formal NMA for adverse events could not be undertaken, which restricts the ability to compare the safety profiles of interventions; this underscores the need for future RCTs to incorporate standardized and comprehensive safety outcome reporting.**

Conclusion

This NMA reinforces the renoprotective roles of ACEIs and ARBs in IgAN, highlighting emerging therapies such as ERAs. Sparsentan, a dual ERA and angiotensin receptor blocker, presents the most excellent efficacy in reducing proteinuria, suggesting its potential as a promising treatment option. However, its superiority over other agents cannot be statistically confirmed. Furthermore, Benazepril, Enalapril, and Irbesartan rank highest among RAAS inhibitors, demonstrating superior proteinuria reduction and eGFR preservation.

330 **Acknowledgments:** None

331 **Funding Source:** None

332 **Author Contributions:** M.E. contributed to the study design and drafting. Y.I. and G.G. worked
333 on the study search, quality check, data extraction, and analysis. A.G., S.V., and C.B. worked on
334 data interpretation and the revision process. All authors have read the manuscript and agree with
335 its content and data.

336 **Data Availability:** The corresponding author shall make the datasets available upon reasonable
337 request.

338 **Ethical Statement:** Institutional Review Board approval was waived due to the nature of the meta-
339 analysis.

340 **Conflict of Interest:** The authors report no conflicts of interest in this work.

341

342 References

- 343 1. Lee M, Suzuki H, Nihei Y, Matsuzaki K, Suzuki Y. Ethnicity and IgA nephropathy:
344 worldwide differences in epidemiology, timing of diagnosis, clinical manifestations, management
345 and prognosis. *Clinical Kidney Journal*. 2023;16(Supplement_2):ii1-ii8. doi:10.1093/ckj/sfad199
- 346 2. Kiryluk K, Freedberg DE, Radhakrishnan J, et al. Global Incidence of IgA Nephropathy
347 by Race and Ethnicity: A Systematic Review. *Kidney360*. 2023;4(8):1112-1122.
348 doi:10.34067/kid.0000000000000165
- 349 3. Kwon CS, Daniele P, Forsythe A, Ngai C. A Systematic Literature Review of the
350 Epidemiology, Health-Related Quality of Life Impact, and Economic Burden of Immunoglobulin
351 A Nephropathy. *J Health Econ Outcomes Res*. 2021;8(2):36-45. doi:10.36469/001c.26129
- 352 4. Cattran DC, Floege J, Coppo R. Evaluating Progression Risk in Patients With
353 Immunoglobulin A Nephropathy. *Kidney International Reports*. 2023/12/01/ 2023;8(12):2515-
354 2528. doi:<https://doi.org/10.1016/j.ekir.2023.09.020>
- 355 5. Bensink ME, Goldschmidt D, Zhou ZY, Wang K, Lieblisch R, Bunke MC. Kidney Failure
356 Attributed to Immunoglobulin A Nephropathy: A USRDS Retrospective Cohort Study of
357 Epidemiology, Treatment Modalities, and Economic Burden. *Kidney Med*. Feb 2024;6(2):100759.
358 doi:10.1016/j.xkme.2023.100759
- 359 6. Lerma EV, Bensink ME, Thakker KM, et al. Impact of Proteinuria and Kidney Function
360 Decline on Health Care Costs and Resource Utilization in Adults With IgA Nephropathy in the
361 United States: A Retrospective Analysis. *Kidney Medicine*.
362 2023;5(9)doi:10.1016/j.xkme.2023.100693
- 363 7. Del Vecchio L, Allinovi M, Comolli S, Peiti S, Rimoldi C, Locatelli F. Drugs in
364 Development to Treat IgA Nephropathy. *Drugs*. May 2024;84(5):503-525. doi:10.1007/s40265-
365 024-02036-1
- 366 8. Bagchi S, Mani K, Swamy A, et al. Supportive Management of IgA Nephropathy With
367 Renin-Angiotensin Blockade, the AIIMS Primary IgA Nephropathy Cohort (APPROACH) Study.
368 *Kidney International Reports*. 2021/06/01/ 2021;6(6):1661-1668.
369 doi:<https://doi.org/10.1016/j.ekir.2021.02.018>
- 370 9. Barratt J, Liew A, Yeo SC, et al. Phase 2 Trial of Cemdisiran in Adult Patients with IgA
371 Nephropathy: A Randomized Controlled Trial. *Clin J Am Soc Nephrol*. Apr 1 2024;19(4):452-462.
372 doi:10.2215/cjn.0000000000000384
- 373 10. Lim RS, Yeo SC, Barratt J, Rizk DV. An Update on Current Therapeutic Options in IgA
374 Nephropathy. *J Clin Med*. Feb 7 2024;13(4)doi:10.3390/jcm13040947
- 375 11. Roccatello D, Careddu A, Ferro M, et al. The steroid-sparing effects of a mycophenolate
376 mofetil-based regimen in the management of immunoglobulin A nephropathy in patients with
377 histologically active lesions: A comparison with a control cohort receiving conventional therapy.
378 *J Nephrol*. Nov 2023;36(8):2223-2231. doi:10.1007/s40620-023-01636-6
- 379 12. Floege J, Bernier-Jean A, Barratt J, Rovin B. Treatment of patients with IgA nephropathy:
380 a call for a new paradigm. *Kidney International*. 2025/01/31/
381 2025;doi:<https://doi.org/10.1016/j.kint.2025.01.014>
- 382 13. Iino Y, Hayashi M, Kawamura T, et al. Interim evidence of the renoprotective effect of the
383 angiotensin II receptor antagonist losartan versus the calcium channel blocker amlodipine in
384 patients with chronic kidney disease and hypertension: a report of the Japanese Losartan Therapy
385 Intended for Global Renal Protection in Hypertensive Patients (JLIGHT) Study. *Clin Exp Nephrol*.
386 Sep 2003;7(3):221-30. doi:10.1007/s10157-003-0241-3

14. Hunter-Dickson M, Wong MG. The role of endothelin receptor antagonists in IgA nephropathy. *Nephrology (Carlton)*. Sep 2024;29 Suppl 2:30-33. doi:10.1111/nep.14364
15. Open Science Framework. Efficacy of Antihypertensive Therapy in Primary IgA Nephropathy in Adults: A Systematic Review and Network Meta-Analysis of Randomized Controlled Trials. . <https://osfio/typj6> (Published 2025, Accessed on March 25, 2025)doi:doi:10.17605/OSF.IO/TYPJ6.
16. Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Bmj*. Mar 29 2021;372:n71. doi:10.1136/bmj.n71
17. Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in randomised trials. *Bmj*. Aug 28 2019;366:l4898. doi:10.1136/bmj.l4898
18. Shao S-C, Kuo L-T, Huang Y-T, Lai P-C, Chi C-C. Using Grading of Recommendations Assessment, Development, and Evaluation (GRADE) to rate the certainty of evidence of study outcomes from systematic reviews: A quick tutorial. *Dermatologica Sinica*. 2023;41(1):3-7. doi:10.4103/ds.DS-D-22-00154
19. Bannister KM, Weaver A, Clarkson AR, Woodroffe AJ. Effect of angiotensin-converting enzyme and calcium channel inhibition on progression of IgA nephropathy. *Contrib Nephrol*. 1995;111:184-92; discussion 192-3. doi:10.1159/000423895
20. Heerspink HJL, Radhakrishnan J, Alpers CE, et al. Sparsentan in patients with IgA nephropathy: a prespecified interim analysis from a randomised, double-blind, active-controlled clinical trial. *Lancet*. May 13 2023;401(10388):1584-1594. doi:10.1016/s0140-6736(23)00569-x
21. Heerspink HJL, Jardine M, Kohan DE, et al. Atrasentan in Patients with IgA Nephropathy. *N Engl J Med*. Feb 6 2025;392(6):544-554. doi:10.1056/NEJMoa2409415
22. Li PK, Leung CB, Chow KM, et al. Hong Kong study using valsartan in IgA nephropathy (HKVIN): a double-blind, randomized, placebo-controlled study. *Am J Kidney Dis*. May 2006;47(5):751-60. doi:10.1053/j.ajkd.2006.01.017
23. Maschio G, Cagnoli L, Claroni F, et al. ACE inhibition reduces proteinuria in normotensive patients with IgA nephropathy: a multicentre, randomized, placebo-controlled study. *Nephrol Dial Transplant*. 1994;9(3):265-9.
24. Nakamura T, Ushiyama C, Suzuki S, et al. Effects of angiotensin-converting enzyme inhibitor, angiotensin II receptor antagonist and calcium antagonist on urinary podocytes in patients with IgA nephropathy. *Am J Nephrol*. Sep-Oct 2000;20(5):373-9. doi:10.1159/000013619
25. Park HC, Xu ZG, Choi S, et al. Effect of losartan and amlodipine on proteinuria and transforming growth factor-beta1 in patients with IgA nephropathy. *Nephrol Dial Transplant*. Jun 2003;18(6):1115-21. doi:10.1093/ndt/gfg090
26. Praga M, Gutiérrez E, González E, Morales E, Hernández E. Treatment of IgA nephropathy with ACE inhibitors: a randomized and controlled trial. *J Am Soc Nephrol*. Jun 2003;14(6):1578-83. doi:10.1097/01.asn.0000068460.37369.dc
27. Remuzzi A, Perico N, Sangalli F, et al. ACE inhibition and ANG II receptor blockade improve glomerular size-selectivity in IgA nephropathy. *Am J Physiol*. Mar 1999;276(3):F457-66. doi:10.1152/ajprenal.1999.276.3.F457
28. Shi Y, Chen W, Jalal D, et al. Clinical outcome of hyperuricemia in IgA nephropathy: a retrospective cohort study and randomized controlled trial. *Kidney Blood Press Res*. 2012;35(3):153-60. doi:10.1159/000331453
29. Shimizu A, Takei T, Uchida K, Tsuchiya K, Nitta K. Low-dose losartan therapy reduces proteinuria in normotensive patients with immunoglobulin A nephropathy. *Hypertens Res*. Sep 2008;31(9):1711-7. doi:10.1291/hypres.31.1711

30. Szeto CC, Kwan BC, Chow KM, Leung CB, Li PK. The safety and short-term efficacy of aliskiren in the treatment of immunoglobulin a nephropathy--a randomized cross-over study. *PLoS One*. 2013;8(5):e62736. doi:10.1371/journal.pone.0062736
31. Woo KT, Chan CM, Choong HL, et al. High dose Losartan and ACE gene polymorphism in IgA nephritis. *Genomic Med*. Dec 2008;2(3-4):83-91. doi:10.1007/s11568-009-9030-8
32. Noor SM, Abuazzam F, Mathew R, Zhang Z, Abdipour A, Norouzi S. IgA nephropathy: a review of existing and emerging therapies. *Front Nephrol*. 2023;3:1175088. doi:10.3389/fneph.2023.1175088
33. Rovin BH, Adler SG, Barratt J, et al. KDIGO 2021 Clinical Practice Guideline for the Management of Glomerular Diseases. *Kidney International*. 2021;100(4):S1-S276. doi:10.1016/j.kint.2021.05.021
34. Vivarelli M, Samuel S, Coppo R, et al. IPNA clinical practice recommendations for the diagnosis and management of children with IgA nephropathy and IgA vasculitis nephritis. *Pediatr Nephrol*. Feb 2025;40(2):533-569. doi:10.1007/s00467-024-06502-6
35. Maixnerova D, Hartinger J, Tesar V. Expanding options of supportive care in IgA nephropathy. *Clin Kidney J*. Dec 2023;16(Suppl 2):ii47-ii54. doi:10.1093/ckj/sfad201
36. Alcocer LA, Bryce A, De Padua Brasil D, et al. The Pivotal Role of Angiotensin-Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers in Hypertension Management and Cardiovascular and Renal Protection: A Critical Appraisal and Comparison of International Guidelines. *Am J Cardiovasc Drugs*. Nov 2023;23(6):663-682. doi:10.1007/s40256-023-00605-5
37. Bomback AS, Toto R. Dual Blockade of the Renin-Angiotensin-Aldosterone System: Beyond the ACE Inhibitor and Angiotensin-II Receptor Blocker Combination. *American Journal of Hypertension*. 2009;22(10):1032-1040. doi:10.1038/ajh.2009.138
38. Kohan DE, Bedard PW, Jenkinson C, Hendry B, Komers R. Mechanism of protective actions of sparsentan in the kidney: lessons from studies in models of chronic kidney disease. *Clin Sci (Lond)*. Jun 5 2024;138(11):645-662. doi:10.1042/cs20240249
39. Bailey CJ, Day C, Bellary S. Renal Protection with SGLT2 Inhibitors: Effects in Acute and Chronic Kidney Disease. *Curr Diab Rep*. Jan 2022;22(1):39-52. doi:10.1007/s11892-021-01442-z
40. Lv J, Wong MG, Hladunewich MA, et al. Effect of Oral Methylprednisolone on Decline in Kidney Function or Kidney Failure in Patients With IgA Nephropathy: The TESTING Randomized Clinical Trial. *JAMA*. 2022;327(19):1888-1898. doi:10.1001/jama.2022.5368

Table 1: Characteristics of Included Studies

Study	Country	Total	Age (years)	Male (%)	Scr (mg/dL)	eGFR ^a	Add-on Therapy	Follow-up	Inclusion Criteria	Therapy
Bannister 1995	Australia	23	51	17 (74%)	N. S	67.8 (8.4)	No	12m	IgAN with eGFR 30–90, with hp	Nifedipine, Enalapril
Heerspink 2023	Intl	404	46	282 (70%)	N. S	57 (24)	45%	110w	IgAN with proteinuria > 1 g/day, eGFR ≥ 30	Sparsentan 400 mg daily, Irbesartan 300 mg daily
Heerspink 2025	Intl	270	45	159 (59%)	N. S	59 (24)	99%	132w	IgAN with proteinuria > 1 g/day, Scr < 2.8 mg/dL	Atrasentan 0.75 mg daily, Placebo
Li 2006	Hong Kong	109	41	30 (28%)	1.2 (0.5)	83 (37)	61%	104w	IgAN with proteinuria > 1 g/day, Scr < 1.4 mg/dL, without hp	Placebo, Valsartan 80 mg daily
Maschio 1994	Italy	39	33	26 (67%)	1.0 (0.2)	103 (23)	No	4m	IgAN with proteinuria ≤ 3 g/day, Scr > 80 mg/dL, without hp	Placebo, Fosinopril 20 mg daily
Nakamura 2000	Japan	32	33	14 (44%)	0.8 (0.2)	111 (13)	No	3m	IgAN with proteinuria > 1 g/day, Scr < 3.0 mg/dL	Verapamil 120 mg daily, Trandolapril 2 mg daily, Candesartan 8 mg daily, Placebo
Park 2003	Korea	36	42	18 (50%)	1.5 (0.6)	63 (23)	50%	12w	IgAN with proteinuria > 0.5 g/day, Scr ≤ 1.5 mg/dL	Losartan 50 mg daily, Amlodipine 5 mg daily
Praga 2003	Spain	44	29	27 (61%)	1 (0.2)	101 (23)	45%	76m	IgAN with proteinuria 0.5–4 g/day, Scr 0.9–2.4 mg/dL	Enalapril 5–40 mg daily, Usual care
Remuzzi 1999	Italy	20	20–65	4 (20%)	N. S	60 (18)	40%	4w	IgAN with proteinuria > 0.5 g/day, Scr < 4 mg/dL	Enalapril 20 mg daily, Irbesartan 100 mg daily
Shi 2002	China	131	29	87 (66%)	1.3 (0.8)	N. S	48%	18m	IgAN with proteinuria > 0.4 g/day, without hp	Benazepril 10–20 mg daily, Usual care
Shimizu 2008	Japan	36	36	17 (47%)	1 (0.2)	74 (17)	No	12m	IgAN with proteinuria > 1 g/day, with hp	Losartan 12.5 mg daily, Usual care
Szeto 2013	Hong Kong	21	45	7 (33%)	1.5 (0.7)	58 (29)	100%	16w	IgAN with proteinuria > 1 g/day, Scr > 1.6 mg/dL	Aliskiren 300 mg daily, Placebo
Woo 2008	Singapore	207	N. S	97 (47%)	N. S	62 (21)	50%	6y	IgAN with eGFR 30–90, with hp	Losartan 100–200 mg daily, Enalapril 10–20 mg daily

Intl: international, Hp: hypertension; Scr: serum creatinine; N.S.: not specified; m: month; w: week; y: year;

Table 2: Network Meta-Analysis of Antihypertensive Agents on Proteinuria

Sparsentan												
-0.4 [-2.8; 1.9]	Benazepril											
-0.7 [-1.7; 0.3]	-0.3 [-2.4; 1.8]	Irbesartan										
-0.9 [-2.9; 0.9]	-0.5 [-1.9; 0.9]	-0.2 [-1.8; 1.4]	Enalapril									
-1.0 [-3.1; 1.0]	-0.6 [-1.9; 0.7]	-0.3 [-2.1; 1.5]	-0.1 [-0.9; 0.7]	Losartan								
-1.6 [-3.9; 0.8]	-1.1 [-2.7; 0.5]	-0.8 [-3.0; 1.3]	-0.6 [-2.0; 0.8]	-0.5 [-1.9; 0.8]	Trandolapril							
-1.6 [-3.9; 0.8]	-1.1 [-2.7; 0.5]	-0.8 [-3.0; 1.3]	-0.6 [-2.0; 0.8]	-0.5 [-1.9; 0.8]	-0.0 [-1.2; 1.2]	Candesartan						
-1.7 [-4.3; 0.9]	-1.3 [-3.2; 0.6]	-1.0 [-3.4; 1.4]	-0.8 [-2.5; 1.0]	-0.7 [-2.4; 1.0]	-0.1 [-1.7; 1.4]	-0.1 [-1.7; 1.4]	Atrasentan					
-1.9 [-4.5; 0.8]	-1.4 [-3.5; 0.6]	-1.1 [-3.6; 1.4]	-0.9 [-2.8; 1.0]	-0.8 [-2.7; 1.0]	-0.3 [-2.0; 1.4]	-0.3 [-2.0; 1.4]	-0.1 [-1.8; 1.5]	Aliskiren				
-1.9 [-4.6; 0.7]	-1.5 [-3.5; 0.5]	-1.2 [-3.7; 1.2]	-1.0 [-2.8; 0.8]	-0.9 [-2.7; 0.9]	-0.4 [-2.0; 1.3]	-0.4 [-2.0; 1.3]	-0.2 [-1.8; 1.3]	-0.1 [-1.8; 1.6]	Fosinopril			
-1.9 [-3.9; 0.2]	-1.4 [-2.5; -0.3]	-1.1 [-2.9; 0.7]	-0.9 [-1.7; -0.1]	-0.8 [-1.6; -0.1]	-0.3 [-1.4; 0.8]	-0.3 [-1.5; 0.9]	-0.1 [-1.7; 1.4]	0.0 [-1.7; 1.7]	0.1 [-1.6; 1.7]	Usual care		
-2.2 [-4.8; 0.5]	-1.7 [-3.7; 0.2]	-1.4 [-3.9; 1.0]	-1.2 [-3.0; 0.6]	-1.1 [-2.9; 0.6]	-0.6 [-2.2; 1.0]	-0.6 [-2.3; 1.0]	-0.5 [-2.0; 1.1]	-0.3 [-2.0; 1.4]	-0.2 [-1.9; 1.4]	-0.3 [-1.9; 1.3]	Valsartan	
-2.4 [-4.7; 0.0]	-1.9 [-3.5; -0.3]	-1.6 [-3.8; 0.5]	-1.4 [-2.8; -0.0]	-1.3 [-2.7; 0.0]	-0.8 [-2.0; 0.4]	-0.8 [-2.0; 0.4]	-0.7 [-1.7; 0.4]	-0.5 [-1.8; 0.8]	-0.4 [-1.6; 0.8]	-0.5 [-1.7; 0.7]	-0.2 [-1.3; 0.9]	Placebo


Table 3: Network Meta-Analysis of Antihypertensive Agents on eGFR

Sparsentan										
4.4 [-16.4; 25.2]	Irbesartan									
6.4 [-26.9; 39.7]	2.0 [-24.0; 28.0]	Enalapril								
8.6 [-35.2; 52.3]	4.2 [-34.3; 42.6]	2.1 [-26.1; 30.4]	Trandolapril							
8.8 [-28.1; 45.8]	4.4 [-26.1; 34.9]	2.4 [-13.5; 18.3]	0.3 [-28.2; 28.8]	Losartan						
12.6 [-37.1; 62.2]	8.2 [-36.9; 53.2]	6.2 [-30.6; 42.9]	4.0 [-30.4; 38.4]	3.7 [-33.2; 40.7]	Valsartan					
12.6 [-30.7; 55.8]	8.2 [-29.8; 46.0]	6.2 [-21.4; 33.7]	4.0 [-20.3; 28.3]	3.7 [-24.1; 31.5]	0.0 [-24.3; 24.3]	Placebo				
12.6 [-31.2; 56.3]	8.2 [-30.3; 46.6]	6.2 [-22.1; 34.4]	4.0 [-21.1; 29.1]	3.7 [-24.8; 32.2]	0.0 [-34.4; 34.4]	0.0 [-24.3; 24.3]	Candesartan			
14.6 [-35.4; 64.5]	10.2 [-35.3; 55.6]	8.2 [-29.1; 45.4]	6.0 [-28.9; 40.9]	5.7 [-31.7; 43.2]	2.0 [-32.9; 36.9]	2.0 [-23.1; 27.1]	2.0 [-32.9; 36.9]	Fosinopril		
14.6 [-21.6; 50.7]	10.2 [-19.4; 39.7]	8.2 [-5.9; 22.2]	6.0 [-18.6; 30.6]	5.7 [-8.7; 20.2]	2.0 [-32.0; 36.0]	2.0 [-21.7; 25.7]	2.0 [-22.6; 26.6]	0.0 [-34.5; 34.5]	Usual Care	
20.9 [-33.9; 75.8]	16.6 [-34.2; 67.3]	14.6 [-29.1; 58.1]	12.4 [-29.2; 54.0]	12.1 [-31.6; 55.9]	8.4 [-33.2; 50.0]	8.4 [-25.4; 42.2]	8.4 [-33.2; 50.0]	6.4 [-35.7; 48.5]	6.4 [-34.9; 47.7]	Aliskiren

Table 4: Network Meta-Analysis of Antihypertensive Agents on Serum Creatinine

Enalapril						
-0.16 [-1.03; 0.71]	Benazepril					
-0.72 [-1.53; 0.10]	-0.56 [-0.95; -0.17]	Losartan				
-0.80 [-1.59; -0.01]	-0.64 [-0.99; -0.29]	-0.08 [-0.26; 0.10]	Usual care			
-0.90 [-1.73; -0.07]	-0.74 [-1.16; -0.32]	-0.18 [-0.49; 0.12]	-0.10 [-0.34; 0.14]	Trandolapril		
-0.90 [-1.73; -0.07]	-0.74 [-1.16; -0.32]	-0.18 [-0.47; 0.11]	-0.10 [-0.33; 0.13]	0.00 [-0.27; 0.27]	Placebo	
-1.00 [-1.82; -0.18]	-0.84 [-1.24; -0.44]	-0.28 [-0.55; -0.01]	-0.20 [-0.40; -0.00]	-0.10 [-0.34; 0.14]	-0.10 [-0.33; 0.13]	Candesartan

Figure 1: Direct Comparisons of Proteinuria Reduction Across Treatments



A: Proteinuria, B: Estimated Glomerular Filtration Rate, C: Serum Creatinine