

Type of Article: Case Report

Title: Occult Primary Osteosarcoma of the Rib in an Adult Male: A Case Report

Authors: Gervie Flor Batucan¹, Mark Ando¹

¹: Department of Internal Medicine, Southwestern University Medical Center, Cebu City,
Philippines

Corresponding: Gervie Flor Batucan

Address: Urgello St. Sambag 2, Cebu City, Philippines, Department of Internal Medicine,
Southwestern University Medical Center

Email: gervieflor@gmail.com

Tel: +63 998 539 6761

Clinical Question Box

What is the appropriate approach for adults with primary osteosarcoma of the rib?

Primary osteosarcoma of the rib is rare; however, management generally adheres to standard osteosarcoma protocols. The recommended approach includes wide surgical resection followed by adjuvant chemotherapy. Evidence indicates that patients who undergo complete resection can achieve a 5-year survival rate of up to 80%, underscoring the importance of early diagnosis and comprehensive treatment.

Abstract

Background: Osteosarcoma is the most common primary bone malignancy in children and young adults, although it rarely occurs in flat bones such as the ribs. **Case**

Presentation: A 31-year-old asymptomatic man was incidentally found to have a homogeneous density on the right anterior chest wall during a pre-employment chest radiograph. Chest CT scan revealed an enhancing mass measuring $10.6 \times 6.5 \times 4.6$ cm, with osseous involvement of the third thoracic vertebral body as well as the right third and fourth ribs. Biopsy confirmed a low-grade fibroblastic osteosarcoma. The patient underwent exploratory thoracotomy, en bloc resection of the second to fifth right ribs, and chest wall reconstruction. He subsequently completed six cycles of adjuvant chemotherapy with cisplatin (100 mg/m^2) and doxorubicin (75 mg/m^2) every 21 days, experiencing only grade 1 adverse events. **Conclusion:** This case highlights the importance of identifying incidental findings, making timely diagnoses, and providing comprehensive treatment to optimize outcomes for rare presentations of osteosarcoma.

Keywords: Osteosarcoma, Doxorubicin, Cisplatin, case report

Introduction

Osteosarcoma is the most common primary bone malignancy, particularly affecting children and adolescents.¹ Approximately 54.2% of cases are diagnosed in individuals aged 10 to 24 years, which coincides with periods of rapid growth, while about 9.6% are found in those over 60.² Males are more commonly affected than females, with an overall male-to-female ratio of 1.3:1. Osteosarcoma that occurs in the context of prior radiation and chemotherapy is classified as a secondary (or subsequent) osteosarcoma.⁴ A hospital-based review in the Philippines identified osteosarcoma as the most prevalent bone tumor, followed by plasma cell myeloma, chondrosarcoma, giant cell tumor, and Ewing's sarcoma.⁵

Bone pain is the primary symptom of osteosarcoma, often beginning during physical activity and progressing to occur even at rest. A prior traumatic event may be reported, although this is not always present.⁶ Systemic signs such as weight loss, fever, fatigue, and general malaise can occur, although their absence does not exclude osteosarcoma.⁷ The femur, tibia, and humerus are the most frequently affected sites, while less common locations include the skull, jaw, and pelvis.⁸ A case was observed in which the patient exhibited no signs or symptoms of bone cancer, despite its occurrence in an

57 uncommon site.

Just Accepted

Case Presentation

A 31-year-old man underwent a routine pre-employment chest X-ray, which revealed a mass on the right anterior chest wall (Figure 1). He was asymptomatic, and his medical and personal history were unremarkable. On physical examination, breath sounds were decreased in the right middle lung field, with no crackles or wheezes. The left lung field was normal. Further diagnostic workup was recommended. Chest CT scan with contrast revealed bony lesions in the right third and fourth ribs, with an associated soft tissue mass measuring approximately $10.6 \times 6.5 \times 4.6$ cm, which enhanced markedly post-contrast. These findings were suspicious for malignancy. Core needle biopsy of the mass indicated a low-grade fibroblastic osteosarcoma (Figure 2). A subsequent PET-CT scan revealed no evidence of distant metastasis, and the tumor was classified as stage IB (T2N0M0). Given the early stage of disease and low-grade features on pathological evaluation, neoadjuvant chemotherapy was not pursued. The patient subsequently underwent exploratory thoracotomy with resection of the right second to fifth ribs, en bloc tumor excision, chest wall reconstruction, and chest tube placement (Figure 3). The postoperative course was uneventful, and he was discharged on postoperative day 7.

At one-month follow up, chest X-ray (Figure 1B) and CT scan (Figure 1C)

showed interval resection of the fourth rib mass with metallic prostheses replacing the right fourth and fifth ribs. The mass at the vertebral end of the third rib remained unchanged. Imaging also showed decreased right lung volume, elevated right hemidiaphragm, and mild pleurodiaphragmatic adhesion. The patient received six cycles of adjuvant chemotherapy with Cisplatin (100 mg/m²) and Doxorubicin (75 mg/m²) every 21 days. Only grade 1 adverse effects were observed, and treatment was completed successfully. A follow-up chest CT scan one month after chemotherapy completion showed no recurrence, as well as stable placement of rib implants.

The patient remains asymptomatic and in good condition, expressing satisfaction with the successful tumor resection and completion of chemotherapy. Follow-up visits are scheduled every 3 months, with continued surveillance planned for a total duration of 5 years.

Discussion

This case illustrates an incidental finding of low-grade fibroblastic osteosarcoma in an asymptomatic adult male, which accounts for only 1.25% of all osteosarcoma cases.⁹ Diagnosis was confirmed through imaging and biopsy, followed by successful en bloc resection and adjuvant chemotherapy. The patient recovered fully, with no recurrence on follow up. This highlights the importance of early detection and comprehensive management even in atypical presentations.

A PubMed search using the keywords “osteosarcoma” and “rib” yielded a summary of the clinical and pathological characteristics of 67 patients diagnosed with rib osteosarcoma, including the present case. The mean age was 29.3 years (standard deviation: 16.7), with a male-to-female ratio of 1.31:1 (38 males, 29 females). Primary osteosarcoma accounted for 87.3% (48/55) of cases, with tumors most commonly located in the first to sixth ribs or overlapping rib regions (57.8%, 37/64). The average maximum tumor size was 7.8 cm (standard deviation: 3.8). Interestingly, 37.9% (25/66) of patients were asymptomatic at diagnosis. Metastatic disease was reported in 11.9% (8/67). High-grade histologic subtypes were identified in 95.9% (47/49) of cases. Surgical resection and chemotherapy were each performed in 90.9% (60/66) of patients, with favorable

outcomes achieved in 56.3% (36/64).

Low-grade intraosseous osteosarcoma is a rare subtype of osteosarcoma that comprised 5.1% of cases in this literature review. It is characterized by osteoid or immature bone formation. For rib and other chest wall tumors, wide surgical excision is the primary treatment modality that ensures clear margins and reduces recurrence. When large sections are removed, reconstruction is required to maintain respiratory function. Complete resection of the primary tumor and any metastases is critical for a potential cure, emphasizing the importance of preserving function.

Chemotherapy, often administered before and after the surgery, typically includes doxorubicin and cisplatin due to their effectiveness and tolerability, as recommended by the NCCN.¹⁴ While the 5-year survival rates for rib osteosarcoma remain modest, strategies such as dose intensification are under investigation. Adverse effects may include myelosuppression, renal dysfunction, mucositis, and cardiotoxicity, although not all patients are affected.¹⁵ Long-term follow up is essential, with studies like EURAMOS-1 showing favorable outcomes in non-metastatic cases after complete resection.¹⁶ The poorer prognosis of this condition is linked to factors such as older age, male sex, large tumor size, axial location, poor chemotherapy response, or absence of surgery.¹⁷

This case report has several limitations. First, the follow-up duration is relatively short, which limits the evaluation of long-term oncologic outcomes, such as recurrence or survival. Second, functional outcomes, including postoperative pulmonary function and quality of life, were not formally assessed, which may reduce the clinical applicability of the findings. Additionally, advanced reconstruction techniques, such as 3D printing, were not available due to institutional limitations; therefore, a metallic rib prosthesis was selected based on its availability and suitability.

Conclusion

This case highlights a rare, asymptomatic presentation of low-grade osteosarcoma in the rib, discovered incidentally on routine imaging. Management with en bloc resection, reconstruction, and adjuvant chemotherapy led to a favorable outcome. Rib osteosarcoma remains uncommon and often lacks specific symptoms. This fact underscores the importance of early detection, complete surgical removal, and ongoing surveillance for optimal prognosis.

Acknowledgment

We appreciate the patient's cooperation in data collection.

Funding Source

No financial support was sought from any source.

Author Contributions

G.B. was responsible for data curation, data interpretation, and drafting of the original manuscript. G.B. and M.A. made substantial contributions to revising the manuscript draft. All authors have read and approved the manuscript and agree with its content and data.

Data Availability

The datasets used in the current study can be obtained from the corresponding author upon reasonable request.

Ethical Statement

This article does not involve the participation of any animals. The patient provided written informed consent for the publication of this report and accompanying images.

Conflict of Interest

The authors report no conflicts of interest in this work.


Table 1: Clinical and Pathological Characteristics of Patients with Rib Osteosarcoma

Items	Value ^a
Number of Cases (n)	67
Age (years)	29.3 (16.7)
Sex (Male-to-Female Ratio)	38/29 (1.31:1)
Primary Osteosarcoma	48/55 (87.3%)
Location (First to Sixth Rib or Overlapping Sites)	37/64 (57.8%)
Maximum Tumor Diameter (cm)	7.8 (3.8)
Asymptomatic Presentation	25/66 (37.9%)
Presence of Metastasis	8/67 (11.9%)
Histologic Classification (High-grade) ^b	47/49 (95.9%)
Surgical Intervention	60/66 (90.9%)
Chemotherapy Administered	60/66 (90.9%)
Favorable Outcome Achieved	36/64 (56.3%)

a: Data are presented as mean and standard deviation for continuous variables.


b: The remaining two cases were classified as low-grade.

159 Figure 1. Radiograph Posteroanterior View showing mass on the right anterior chest wall



160
161 A: Chest Radiograph, pre-operative; B: Chest Radiograph, post-operative; C: Chest Computed
162 tomography, post-operative
163

164 Figure 2. Pathological Diagnosis Based on H&E-Stained Sections



165

166

Just Acc

167 Figure 3. Intraoperative chest wall reconstruction



168

169

Just Accepted