

Meta-Analysis

Clinical Features of Pneumocystis Pneumonia in Non-Human Immunodeficiency Virus-Infected Patients: A Systemic Review and Meta-Analysis

Masashi Ishihara, Shigeru Tanzawa, Takeshi Honda, Yasuko Ichikawa, Kiyotaka Watanabe, Nobuhiko Seki*

Division of Oncology, Department of Internal Medicine, Teikyo University School of Medicine, Tokyo, Japan.

***Corresponding Author:** e-mail: nseki@med.teikyo-u.ac.jp

Submitted: August 06, 2024 **Published:** September 02, 2024

Clinical Question Box

What are the clinical features of pneumocystis pneumonia in non-human immunodeficiency virus-infected patients?

Pneumocystis pneumonia was more likely to occur in men than in women at the age of 60, with a sex ratio of about 3 to 2 (males to females). The main symptoms were dyspnea and fever, which were observed in >70% of patients, accompanied by cough in >50% of patients. Ground glass opacity might offer a clue to diagnosing pneumocystis pneumonia in high-risk patients.

Abstract

Objectives: Pneumocystis pneumonia (PCP) is an opportunistic disease that causes potentially fatal pneumonia in immunocompromised individuals. The clinical features of PCP without HIV remain incompletely understood. **Methods:** This study aimed to identify the clinical features of PCP without HIV in a systematic review following a meta-analysis. **Results:** 65 articles that included 10,133 PCP patients without HIV infection were enrolled. PCP occurred most commonly at age 59.2 years (95% CI: 57.7, 60.7) in a gender ratio of approximately 3 to 2 (males to females). Dyspnea, fever, cough, and sputum were nonspecific clinical findings in 73% (95% CI: 69, 79), 73% (95% CI: 65, 81), 56% (95% CI: 48%, 64%), and 32% (96%CI: 16, 48) of patients, respectively. Viral, bacterial, and fungal co-infection were observed in 28% (95% CI: 13, 44), 19% (95% CI: 13, 25), and 11% (95% CI: 6, 16) of patients, respectively. Laboratory data showed a trend of elevated WBC, LDH, CRP, β -D glucan, and KL-6. Ground glass opacity (GGO) was found in 87% (95% CI: 83, 91) of patients. In-hospital mortality was 41% (95% CI: 35, 46). **Conclusions:** PCP is a life-threatening disease in immune-compromised patients. Despite being a nonspecific clinical finding, GGO might offer a clue to diagnosing PCP in high-risk patients.

Keywords: Pneumocystis pneumonia, clinical features, non-HIV, ground glass opacity.

Introduction

In individuals who are immunocompromised, the opportunistic fungal pathogen *Pneumocystis jirovecii* is capable of causing life-threatening pneumonia as the causative agent of pneumocystis pneumonia (PCP).¹ Human immunodeficiency virus (HIV)-infected patients are at the highest risk of PCP; however, non-HIV-infected immunocompromised patients, including those with chronic corticosteroid use, hematological or solid malignancies, organ transplantation, and immunomodulatory or biological therapy have a substantial risk of developing PCP.^{2,3}

HIV-infected patients usually develop a subacute course of the disease, whereas respiratory insufficiency is usually more severe in non-HIV patients than in the HIV-infected population. *Pneumocystis* is more challenging to detect in non-HIV patients because of the smaller numbers of organisms in the lungs.⁴ In addition, the mortality rate is 30%–50% in patients without HIV and 10%–12% in patients with HIV.⁵ The diagnosis of PCP relies on a critical evaluation of clinical symptoms, risk factors, radiologic features, and microbiological tests. Computed tomography (CT) examination is helpful if there is clinical suspicion of PCP.

A clear overview of PCP in non-HIV patients is essential for diagnostic progress and treatment in clinical practice; however, this is currently lacking, and no meta-analysis has appraised the clinical features in this group. Therefore, we conducted a systematic review and meta-analysis to explore the general characteristics, comorbidity, co-infection, and laboratory and radiographic findings of PCP obtained in observational studies of non-HIV patients.

Methods

Study Overview

The protocol of this systematic review and meta-analysis followed the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. It was registered on the website of the University Hospital Medical Information Network Clinical Trials Registration (UMIN 000046230).^{6,7} Because of the nature of the study, Institutional Review Board approval was not required.

Study Search

Four major online databases (PubMed, Web of Science, Cochrane, and Embase) were searched from January 1, 2000 to October 30, 2022. The following search strategy was used for PubMed: (((((pneumocystis pneumonia) OR (pneumocystis jirovecii)) OR (pneumocystis infections)) OR (pneumocystis carinii)) AND (((non-HIV) OR (non-AIDS)) OR (non-HIV-infected)) OR (HIV-uninfected)) OR (HIV-negative))) AND (((symptom) OR (clinical feature)) OR (diagnosis)). Two authors (H.C. and M.I.) independently screened the titles and abstracts and carefully evaluated the full text to select eligible articles; in cases of discrepancy, a consensus was reached through discussion. Review articles and the included original articles were hand-searched (H.C. and M.I.) for additional research papers that met the inclusion criteria.

Inclusion and Exclusion Criteria

Full-length articles published in any language that provided data regarding the clinical features of PCP were included. The inclusion criteria were: (1) patients with PCP, (2) patients without HIV, and (3) age ≥ 16 years. The exclusion criteria were: (1) detailed data not available, (2) fewer than 10 cases, and (3) limited information regarding clinical features.

Risk of Bias

Two reviewers independently assessed the methodological quality of the selected studies using the Newcastle–Ottawa quality assessment to evaluate the quality of observational studies.⁸ Due to the nature of single-arm studies, the “selection of the non-exposed cohort” and “comparability” domains were not applicable in this study. The modified Newcastle–Ottawa quality assessment scored six stars. Any disagreement between the reviewers was discussed, and agreement was reached by consensus.⁹

Outcomes

The clinical features of PCP were appraised in terms of the following five aspects: general characteristics, laboratory findings, comorbidity, existence of other microorganisms, and radiographic findings. General characteristics included age, sex, body mass index (BMI), prophylaxis, time from symptom onset to treatment, and hospital mortality. The following laboratory findings were collected: white blood cells (WBC), lactate dehydrogenase (LDH), albumin, c-reactive protein (CRP), β -D glucan, and sialylated carbohydrate antigen 6 (KL-6). Comorbidities included hematologic malignancies, solid tumors, immunosuppression (immunosuppressive drugs for nonmalignant diseases), and any transplant. Symptoms included dyspnea, fever, cough, and sputum. Although it is difficult to distinguish between co-infection and detection of the existence of other microorganisms, most previous studies have reported co-infection. Co-infection was classified as bacterial, viral, cytomegalovirus (CMV), and fungal. The radiographic findings were classified as ground glass opacity (GGO), consolidation, lung nodules, septal thickening, pleural effusion, and pneumothorax.

Data Extraction and Statistics

Two authors (M.I. and S.T.) independently extracted the following data from the studies: first author, publication year, country of publication, number of patients with PCP, clinical features, and Newcastle–Ottawa quality assessment-related information. All analyses were performed using Review Manager software version 5.3 (Cochrane Collaboration, Oxford, UK). Figures illustrated using Review Manager were adjusted as necessary. Heterogeneity was evaluated with the I^2 statistic and interpreted as follows: $I^2 = 0\%$, no heterogeneity; $0\% < I^2 < 25\%$, the least heterogeneity; $25\% \leq I^2 < 50\%$, mild heterogeneity; $50\% \leq I^2 < 75\%$, moderate heterogeneity; and $75\% \leq I^2$, strong heterogeneity.¹⁰ A p-value of <0.05 was considered significant.

Results

Study Search and Study Characteristics

A total of 581 articles were identified, including 578 articles by database search and three articles by hand search. Of the 581 articles, 416, 179, and 65 remained after removing duplication, screening for eligibility, and full-length article reading, respectively (Fig. S1). The analysis included 65 retrospective studies (Table 1).^{11–75} All but two were written in English.^{19,32} Of the 65 reports, 15 were from Japan, 13 were from China, 12 were from France, 5 were from Korea and the USA, 3 were from Germany, and 2 were from Israel and Spain. These studies included a total of 10,133 PCP patients without HIV infection. Among the studies, the modified Newcastle–Ottawa scale ranged from 5 to 6 stars. The method for diagnosis of PCP showed some variation among studies (Table S1). Polymerase chain reaction (PCR) was conducted in 43 studies. Clinical and/or radiological findings were also included as diagnostic criteria for PCP in 39 studies. Elevation of β -D-glucan was considered necessary in 6 studies, and no diagnostic criteria were provided in 5 studies.

Table 1. Characteristics of the enrolled studies

Author	Year	Country	Cases	Male	Age	Prophylaxis	Mortality	NOS
Ainoda	2012	Japan	24	10	66.5	ND	45.8% [‡]	6
Asai	2017	Japan	38	20		ND	34.2%	6
Assal	2021	France	133	100	65.0	ND	24.1% [†]	6
Azoulay	2009	France	39			ND	NA	6
Bienvenu	2016	France	461	288		27	20.6% [‡]	6
Bollee	2007	France	56	36	49.3	ND	19.6% [‡]	6
Burghi	2021	France	321	193	57.7	55	24.3% [†]	5

(Continued)

Table 1. Continued

Author	Year	Country	Cases	Male	Age	Prophylaxis	Mortality	NOS
Calero-Bernal	2016	Spain	128	86		7	28.9%	6
Cerón	2014	Chile	45	42		ND	33.0%	6
Choi	2018	Korea	81	51	60.8	ND	64.2%	5
Chou	2014	Taiwan	65	34	49.5	ND	44.6%	6
de Boer	2011	Netherlands	21	11	56.3	ND	9.5% [†]	5
Deodhar	2018	India	29			ND	34.5%	6
Enomoto	2010	Japan	17	9	68.0	0	35.3% [†]	6
Festic	2005	USA	30	13		ND	66.7%	6
Fillatre	2014	France	154			ND	59.2% [‡]	5
Gilmartin	2002	USA	33	22	57.8	3	27.3%	6
Grønseth	2021	Norway	297	180	66.3	ND	21.5%	6
Guo	2014	China	46	28	54.7	ND	15.2%	6
Hamada	2020	Japan	74	40	73.2	1	NA	5
Hardak	2012	Israel	58	28	55.9	ND	17.2%	6
Huang	2016	China	92	40	53.1	ND	76.1%	6
Ideguchi	2020	Japan	50	27	66.3	0	NA	5
Inoue	2019	Japan	562	324	62.9	ND	44.0% [‡]	6
Issa	2018	France	112	74	63.8	ND	33.0%	6
Jin	2019	China	126	71	53.7	ND	34.1% [‡]	5
Kanj	2021	USA	2780	1496	60.2	ND	16.0%	6
Kato	2019	Japan	44	25	62.7	ND	NA	6
Kim	2014	Korea	173	116	56.0	6	35.8%	5
Ko	2019	Korea	51	35	52.8	2	74.5%	6
Kofteridis	2014	Greece	62	43	65.2	21	29.0%	6
Korkmaz Ekren	2018	Turkey	45	43	56.7	3	64.4%	6
Kosaka	2017	Japan	77	47	66.1	1	22.1%	5
Kumagai	2019	Japan	61	28	69.0	ND	31.1%	6
Kunihiro	2015	Japan	70	34	56.4	ND	NA	6
Lahmer	2017	Germany	50	29	59.0	ND	66.0%	6
Lee	2019	Korea	362	211	51.0	94	32.6%	6
Lemiale	2013	France	139	79	49.3	5	25.9% [‡]	6
Li	2014	Taiwan	20	9	50.2	ND	60.0%	5
Liu	2019	China	84	71		ND	46.4% [‡]	5
Mansharamani	2000	Canada	33	22	57.8	ND	39.4%	6
Matsumura	2011	Japan	82	51		ND	24.0% [†]	6
Moon	2011	Korea	88	46	55.5	ND	31.8% [†]	6
Mu	2016	China	105	62		ND	34.3%	5
Mundo	2020	USA	28	15	62.1	ND	71.4%	6
Nakamura	2009	Japan	16	9	58.5	ND	43.8%	5
Nakashima	2017	Japan	54	34	72.7	ND	4.2% [†]	6
Pereira-Díaz	2019	Spain	1204	728	58.0	ND	25.5%	6
Roblot	2002	France	103	61	57.4	4	37.9%	5
Roembke	2014	Germany	30	19	56.4	ND	36.7% [†]	6
Roux	2014	France	321			12	23.4%	6

(Continued)

Table 1. Continued

Author	Year	Country	Cases	Male	Age	Prophylaxis	Mortality	NOS
Sheikholeslami	2013	Iran	89	89	66.5	ND	NA	5
Sun	2021	China	24	12	51.3	ND	25.0% [†]	5
Tamai	2014	Japan	29	14	58.7	ND	41.4%	6
Tang	2021	China	40	26	52.3	ND	NA	6
Tasaka	2010	Japan	21	13	50.7	ND	52.4%	6
Toper	2011	France	41		56.0	7	29.3%	5
Vogel	2007	Germany	27	17		0	NA	6
Wang	2017	China	35	24	54.5	ND	NA	5
Weng	2016	China	82	34	52.6	ND	75.6%	6
Wieruszewski	2020	USA	323	202	63.7	8	NA	5
Wu	2021	Taiwan	62	38	59.7	ND	43.5%	6
Yu	2017	China	70	43		ND	37.1%	6
Zahar	2002	France	39	20		5	NA	6
Zalmanovich	2020	Israel	77	49	65.3	5	44.2%	6

Note: †, 30-day mortality; ‡, 60-day mortality; ¶, Ainoda 2012, 90-day mortality; Bienvenu 2016, 14-day mortality; Fillatre 2014 and Lemiale 2013, ICU mortality; ND, not described; NA, not available; NOS, Newcastle–Ottawa Scale.

General Characteristics and Laboratory Findings

Table 2 lists the general characteristics of patients with PCP without HIV. In an analysis of 50 studies, the mean age was 59.2 years (95% confidence interval (CI): 57.7, 60.7; $I^2 = 95\%$; Fig. S2). Males comprised 60% of patients (95% CI: 58, 63; $I^2 = 85\%$; Fig. S3). Mean BMI was 22.1 kg/m² (95% CI: 20.0, 24.1; $I^2 = 0\%$; Fig. S4), and 8% of patients underwent prophylaxis for PCP (95% CI: 5, 11; $I^2 = 92\%$; Fig. S5). Trimethoprim/sulfamethoxazole was used as prophylaxis in 11 studies. However, detailed information should have been provided. The other eight studies (Table S2) did not state the length of prophylaxis. The mean time from symptom onset to treatment was 8.9 days (95% CI: 6.8, 10.9; $I^2 = 94\%$; Fig. S6), and mean in-hospital mortality due to PCP was 41% (95% CI: 35, 46; $I^2 = 95\%$; Fig. S7) in 34 studies.

Table 2. General characteristics and laboratory findings in patients with PCP without HIV

	Studies (n)	Statistical method	Estimated effect	95% CI
Age (years)	50	IV, random	59.2	57.7, 60.7
Sex (male)	60	IV, random	60	58, 63
BMI (kg/m ²)	10	IV, random	22.1	20.0, 24.1
Prophylaxis (%)	18	IV, random	8	5, 11
WBC (/μL)	18	IV, random	8210	7810, 8610
LDH (IU/L)	31	IV, random	496.4	454, 538
Albumin (mg/dL)	19	IV, random	2.86	2.76, 2.97
CRP (mg/dL)	17	IV, random	11.1	9.5, 12.8
(1,3) β-D glucan (pg/mL)	13	IV, random	137.1	95.1, 179.2
KL-6 (U/mL)	7	IV, random	1073	724, 1423
Time from onset (days)	9	IV, random	8.9	6.8, 10.9
Mortality (%)	34	IV, random	41	35, 46

Note: PCP, pneumocystis pneumonia; HIV, human immunodeficiency virus; CI, confidence interval; IV, inverse variance; BMI, body mass index; WBC, white blood cell; LDH, lactate dehydrogenase; CRP, c-reactive protein; KL-6, sialylated carbohydrate antigen 6.

The laboratory findings revealed elevated WBC, LDH, CRP, β -D glucan, and KL-6 and a decreased albumin value in the overall patients. The average values were as follows: WBC, 8,210/ μ L (95% CI: 7,810, 8,610; $I^2 = 52\%$; Fig. S8); LDH, 496 IU/L (95% CI: 454, 538; $I^2 = 94\%$; Fig. S9); albumin, 2.86 mg/dL (95% CI: 2.76, 2.97; $I^2 = 91\%$; Fig. S10); CRP, 11.1 mg/dL (95% CI: 9.5, 12.8; $I^2 = 89\%$; Fig. S11); β -D glucan, 137.1 pg/dL (95% CI: 95.1, 179.2; $I^2 = 100\%$; Fig. S12); and KL-6, 1073 (95% CI: 724, 1423; $I^2 = 91\%$; Fig. S13).

Comorbidity

The frequencies of comorbidities in patients with PCP without HIV infection are listed in Table S1. The most frequent comorbidities were hematologic malignancies and use of immunosuppressive drugs for nonmalignant diseases, followed by transplant and solid tumor, as follows: 32% (95% CI: 27, 36; $I^2 = 97\%$; Fig. S14); 32% (95% CI: 27, 36; $I^2 = 96\%$; Fig. S15), 19% (95% CI: 16, 23; $I^2 = 95\%$; Fig. S16), and 16% (95% CI: 14, 18; $I^2 = 81\%$; Fig. S17).

Symptoms

Table 3 lists the symptoms reported in at least two studies. The most frequently reported symptoms were dyspnea and fever, followed by cough, sputum, hemoptysis and chest pain, as follows: 73% (95% CI: 68, 79; $I^2 = 94\%$; Fig. S18), 73% (95% CI: 65, 81; $I^2 = 97\%$; Fig. S19), 56% (95% CI: 48, 64; $I^2 = 96\%$; Fig. S20), 32% (95% CI: 16, 48; $I^2 = 95\%$; Fig. S21), 9% (95% CI: 2, 16; $I^2 = 49\%$; Fig. S22), and 17% (95% CI: 0, 37; $I^2 = 85\%$; Fig. S23), respectively. The meta-analysis did not include articles that reported limited fatigue and body-weight loss data.

Table 3. Symptoms of PCP in non-HIV patients

Symptoms	Studies (n)	Statistical method	Effect estimate	95% CI
Dyspnea (%)	29	IV, random	73	68, 79
Fever (%)	27	IV, random	73	65, 81
Cough (%)	29	IV, random	56	48, 64
Sputum (%)	7	IV, random	32	16, 48
Hemoptysis (%)	2	IV, random	9	2, 16
Chest pain (%)	2	IV, random	17	0, 37

Note: PCP, pneumocystis pneumonia; HIV, human immunodeficiency virus; CI, confidence interval; IV, inverse variance.

Co-infection

Table 4 lists data regarding the detected microorganisms. In patients with PCP without HIV infection, viral microorganisms were detected most frequently, followed by bacterial and fungal, as follows: 28% (95% CI: 13, 44; $I^2 = 97\%$; Fig. S24), 19% (95% CI: 13, 25; $I^2 = 89\%$; Fig. S25), and 11% (95% CI: 13, 44; $I^2 = 97\%$; Fig. S26), respectively. Cytomegalovirus (CMV) was observed in 26% (95% CI: 16, 36; $I^2 = 97\%$; Fig. S27) of PCP patients. The methods used to identify CMV, viruses, and bacteria are shown in Tables S3, S4, and S5, respectively. A total of 14 studies reported a positive result for CMV, but only eight described the identification method. Among these, CMV antigenemia assay and real-time PCR were used in three studies each. Positive results were found for CMV in 25.5% and 47.3% of antigenemia assay and real-time PCR, respectively. The co-infection of viruses and bacteria was reported in 7 and 15 studies, respectively. Microbiological confirmation was used as the detection method in most of these studies.

Radiographic Findings

Table 5 summarizes the radiographic findings. GGO was most the frequent radiographic finding, followed by septal thickening, consolidation, pleural effusion, lung nodules, and pneumothorax, as follows: 87% (95% CI: 83, 91; $I^2 = 95\%$; Fig. S28), 41% (95% CI: 24, 58; $I^2 = 91\%$; Fig. S29), 29%

(95% CI: 19, 38; $I^2 = 94\%$; Fig. S30), 20% (95% CI: 13, 28; $I^2 = 90\%$; Fig. S31), 14% (95% CI: 7, 21; $I^2 = 91\%$; Fig. S32), and 4% (95% CI: 2, 7; $I^2 = 63\%$; Fig. S33), respectively.

Table 4. Co-infection of PCP in non-HIV patients

Co-infection	Studies (n)	Statistical method	Effect estimate	95% CI
Viral (%)	7	IV, random	28	13, 44
Cytomegalovirus (%)	14	IV, random	26	16, 36
Bacterial (%)	15	IV, random	19	13, 25
Fungal (%)	11	IV, random	11	6, 16

Note: PCP, pneumocystis pneumonia; HIV, human immunodeficiency virus; CI, confidence interval; IV, inverse variance.

Table 5. Radiographic findings of PCP in non-HIV patients

Finding	Studies (n)	Statistical method	Effect estimate	95% CI
GGO (%)	19	IV, random	87	83, 91
Septal thickening (%)	5	IV, random	41	24, 58
Consolidation (%)	13	IV, random	29	19, 38
Pleural effusion (%)	12	IV, random	20	13, 28
Lung nodules (%)	9	IV, random	14	7, 21
Pneumothorax (%)	7	IV, random	4	2, 7

Note: PCP, pneumocystis pneumonia; HIV, human immunodeficiency virus; GGO, ground glass opacity; IV, inverse variance; CI, confidence interval.

Discussion

The general features of PCP, including clinical presentation and laboratory and radiographic findings, were identified after analysis of 10,133 cases of PCP in patients without HIV. PCP was more likely to occur in men than in women at the age of 60 years, with a sex ratio of about 3 to 2 (males to females). The main symptoms were dyspnea and fever, which were observed in >70% of patients, accompanied by cough in >50% of patients. Most studies did not specify the type of cough; however, cough was accompanied by sputum in >30% of patients. The co-infection was frequently observed, and CMV was detected in >25% of patients. The laboratory findings revealed decreased albumin but no apparent weight loss in PCP patients. Elevated WBC, LDH, CRP, β -D glucan, and KL-6 were also observed in these patients. More than 60% of patients had a comorbidity of either hematologic malignancy or immunosuppressive drug use for nonmalignant disease. Eight percent of PCP patients underwent prophylaxis for PCP, and in-hospital mortality was high (~41%), which is in agreement with the results of previous studies.^{3,34,76}

PCP is of increasing importance in non-HIV immunocompromised patients who present with severe respiratory distress and low fungal loads. β -D glucan level was found to be very useful in the diagnosis of PCP, and seven studies included β -D glucan positivity as a condition for the diagnosis of PCP. In contrast, three studies have reported negative β -D glucan in non-HIV patients, ranging from 12% to 27.2%.^{38,52,56} These reports were all from Japan, which suggests possible racial differences or differences in laboratory examination methods between countries. The high rate of CMV positivity was surprising. CMV generally produces an asymptomatic or minimally symptomatic acute illness in immunocompetent patients. Detecting viral DNA/RNA does not necessarily mean clinically relevant disease but may also indicate viral 'dead' material.

Most importantly, CMV reactivation can be found in sick airways, and low levels of CMV can be found in peripheral blood. However, the enrolled articles provided limited information that

would enable differences to be identified between genetic information and CMV infection. The exact proportion of patients with the existence of CMV and how CMV infection affects the clinical progress of PCP in non-HIV patients remains unclear.

GGO was observed in almost 90% of PCP patients. Given the nonspecific nature of the clinical findings, this radiology finding appears to be an essential clue for confirming a diagnosis of PCP, even in asymptomatic high-risk patients. Numerous studies reported less common radiographic findings such as consolidation, pleural effusion, and lung nodules. Combining co-infection and PCP might be necessary to interpret the various image findings. The diagnostic accuracy of radiology in PCP remains unclear, and further studies are required.

PCP is challenging to diagnose. Asymptomatic lung colonization can occur in people with normal immune systems who may unknowingly become reservoirs (asymptomatic carriers) for the spread of pneumocystis to immunocompromised individuals.⁷⁷ A few articles have discussed the relevance of sputum colonization vs. bronchoalveolar lavage sampling and the sensitivities between PCR and microscopy staining.

There were several limitations in this study. First, high heterogeneity was observed in almost all of the analyses performed due to the observational nature of the studies and their differing purposes. Although random effects were considered in data analysis, selection bias was possible. Second, the detection of pathogens is different from the diagnosis of infection. No uniform criteria were included in studies to differentiate between detecting pathogens and diagnosing disease, especially in the prevalence of CMV infection. Thirdly, all of the clinical features of PCP in patients without HIV infection were conducted in a single-arm meta-analysis. There was no comparison between PCP and community-acquired pneumonia or other types of pneumonia. Fourth, a limited number of studies discussed the clinical features of PCP in terms of different comorbidities; therefore, it was impossible to perform a subgroup analysis of clinical features according to comorbidity.

Conclusions

PCP is a life-threatening disease in immune-compromised patients. Men are more susceptible than women, and there is an association with comorbidities. Dyspnea, fever, and cough were nonspecific clinical findings in PCP, and GGO was observed in most patients.

Acknowledgments

None.

Funding Source

None.

Author Contributions

M.I. contributed to the study search, quality check, data extraction, and drafting. As principal investigators, H.K. and S.T. worked on the study search, quality check, data extraction, and analysis. T.H., Y.I., K.W., and N.S. worked on the interpretation of data and the revision process. All the authors have approved the final version of the manuscript for publication.

Data Availability Statement

The datasets used in the current study are available from the corresponding author upon reasonable request.

Ethical Statement

Not applicable.

Conflicts of Interest

The authors declare that no conflicts of interest exist.

Supplemental Information

Supplemental information for this article can be found online at <https://sup.jclinque.com/api/articles/39/download-suppl>.

References

- [1] Gingerich AD, Norris KA, Mousa JJ. Pneumocystis pneumonia: immunity, vaccines, and treatments. *Pathogens*. February 19, 2021;10(2). doi:10.3390/pathogens10020236.
- [2] Shibata S, Kikuchi T. Pneumocystis pneumonia in HIV-1-infected patients. *Respir Investig*. May 2019;57(3):213–219. doi:10.1016/j.resinv.2019.01.009.
- [3] Cillóniz C, Domínguez C, Álvarez-Martínez MJ, et al. Pneumocystis pneumonia in the twenty-first century: HIV-infected versus HIV-uninfected patients. *Expert Rev Anti Infect Ther*. October 2019;17(10):787–801. doi:10.1080/14787210.2019.1671823.
- [4] Tasaka S. Recent advances in the diagnosis and management of pneumocystis pneumonia. *Tuberc Respir Dis (Seoul)*. April 2020;83(2):132–140. doi:10.4046/trd.2020.0015.
- [5] Datta S, Mahal S, Ravat V, et al. Hospitalization outcomes in pneumocystis pneumonia inpatient population: a comparison between HIV and non-HIV patients. *Cureus*. August 1, 2018;10(8):e3082. doi:10.7759/cureus.3082.
- [6] Hospital Medical Information Network. Clinical features of PCP in non-HIV infected patients: A systematic review and meta-analysis. Updated December 1, 2022. Accessed on November 30, 2023. https://upload.umin.ac.jp/cgi-open-bin/ctr_e/ctr_view.cgi?recptno=R000052760.
- [7] Bernardo WM. PRISMA statement and PROSPERO. *Int Braz J Urol*. May–June 2017;43(3):383–384. doi:10.1590/s1677-5538.10.2017.03.02.
- [8] Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol*. September 2010;25(9):603–605. doi:10.1007/s10654-010-9491-z.
- [9] Chen H, Ishihara M, Horita N, et al. Effectiveness of cell-free and concentrated ascites reinfusion therapy in the treatment of malignancy-related ascites: a systematic review and meta-analysis. *Cancers (Basel)*. September 29, 2021;13(19):4873. doi:10.3390/cancers13194873.
- [10] Higgins JP, Thompson SG, Deeks JJ, et al. Measuring inconsistency in meta-analyses. *BMJ*. September 6, 2003;327(7414):557–560. doi:10.1136/bmj.327.7414.557.
- [11] Ainoda Y, Hirai Y, Fujita T, et al. Analysis of clinical features of non-HIV Pneumocystis jirovecii pneumonia. *J Infect Chemother*. October 2012;18(5):722–728. doi:10.1007/s10156-012-0408-5.
- [12] Asai N, Motojima S, Ohkuni Y, et al. Clinical manifestations and prognostic factors of Pneumocystis jirovecii pneumonia without HIV. *Chemotherapy*. 2017;62(6):343–349. doi:10.1159/000477332.
- [13] Assal M, Lambert J, Chow-Chine L, et al. Prognostic impact of early adjunctive corticosteroid therapy in non-HIV oncology or haematology patients with Pneumocystis jirovecii pneumonia: a propensity score analysis. *PLoS One*. 2021;16(4):e0250611. doi:10.1371/journal.pone.0250611.
- [14] Azoulay É, Bergeron A, Chevret S, et al. Polymerase chain reaction for diagnosing pneumocystis pneumonia in non-HIV immunocompromised patients with pulmonary infiltrates. *Chest*. March 2009;135(3):655–661. doi:10.1378/chest.08-1309.
- [15] Bienvenu AL, Traore K, Plekhanova I, et al. Pneumocystis pneumonia suspected cases in 604 non-HIV and HIV patients. *Int J Infect Dis*. May 2016;46(2):11–17. doi:10.1016/j.ijid.2016.03.018.
- [16] Bollee G, Sarfati C, Thiery G, et al. Clinical picture of Pneumocystis jirovecii pneumonia in cancer patients. *Chest*. October 2007;132(4):1305–1310. doi:10.1378/chest.07-0223.
- [17] Burghi G, Biard L, Roux A, et al. Characteristics and outcome according to underlying disease in non-AIDS patients with acute respiratory failure due to Pneumocystis pneumonia. *Eur J Clin Microbiol Infect Dis*. June 2021;40(6):1191–1198. doi:10.1007/s10096-020-04118-w.

- [18] Calero-Bernal ML, Martin-Garrido I, Donazar-Ezcurra M, et al. Intermittent courses of corticosteroids also present a risk for pneumocystis pneumonia in non-HIV patients. *Can Respir J*. 2016;2016(3):2464791. doi:10.1155/2016/2464791.
- [19] Cerón I, Rabagliati R, Langhaus J, et al. [Pneumocystis jiroveci pneumonia: comparative study of cases in HIV-infected patients and immunocompromised non-HIV-infected patients]. *Rev Chilena Infectol*. August 2014;31(4):417–424. Características clínicas, diagnósticas y pronósticas de pacientes con neumonía por *Pneumocystis jiroveci* en individuos infectados por virus de inmunodeficiencia humana e individuos inmunocomprometidos por otra etiología: comparative study of cases in HIV-infected patients and immunocompromised non-HIV-infected patients. doi:10.4067/s0716-10182014000400007.
- [20] Choi JS, Lee SH, Leem AY, et al. Pneumocystis jirovecii pneumonia (PCP) PCR-negative conversion predicts prognosis of HIV-negative patients with PCP and acute respiratory failure. *PLoS One*. 2018;13(10):e0206231. doi:10.1371/journal.pone.0206231.
- [21] Chou CW, Lin FC, Tsai HC, et al. The impact of concomitant pulmonary infection on immune dysregulation in *Pneumocystis jirovecii* pneumonia. *BMC Pulm Med*. November 19, 2014;14(1):182. doi:10.1186/1471-2466-14-182.
- [22] de Boer MGJ, Gelinck LBS, van Zelst BD, et al. beta-D-Glucan and S-adenosylmethionine serum levels for the diagnosis of *Pneumocystis pneumonia* in HIV-negative patients: a prospective study. *J Infect*. January 2011;62(1):93–100. doi:10.1016/j.jinf.2010.10.007.
- [23] Deodhar D, Koshy JM, John M, et al. Clinical profile of pneumocystis jirovecii infection-A comparative study. *J Assoc Phys India*. January 2018;66(1):28–31. <https://japi.org/q2c4e434/clinical-profile-of-pneumocystis-jirovecii-infection-a-comparative-study>.
- [24] Enomoto T, Azuma A, Kohno A, et al. Differences in the clinical characteristics of *Pneumocystis jirovecii* pneumonia in immunocompromised patients with and without HIV infection. *Respirology*. January 2010;15(1):126–131. doi:10.1111/j.1440-1843.2009.01660.x.
- [25] Festic E, Gajic O, Limper AH, et al. Acute respiratory failure due to *Pneumocystis pneumonia* in patients without human immunodeficiency virus infection—Outcome and associated features. *Chest*. August 2005;128(2):573–579. doi:10.1378/chest.128.2.573.
- [26] Fillatre P, Decaux O, Jouneau S, et al. Incidence of *Pneumocystis jiroveci* pneumonia among groups at risk in HIV-negative patients. *Am J Med*. December 2014;127(12):1242. e11. doi:10.1016/j.amjmed.2014.07.010.
- [27] Gilmartin GS, Koziel H. *Pneumocystis carinii* pneumonia in adult non-HIV disorders. *J Intens Care Med*. November–December 2002;17(6):283–301. doi:10.1177/0885066602238031.
- [28] Grønseth S, Rogne T, Hannula R, et al. Epidemiological and clinical characteristics of immunocompromised patients infected with *Pneumocystis jirovecii* in a twelve-year retrospective study from Norway. *BMC Infect Dis*. July 7, 2021;21(1):659. doi:10.1186/s12879-021-06144-1.
- [29] Guo F, Chen Y, Yang SL, et al. *Pneumocystis pneumonia* in HIV-infected and immunocompromised non-HIV infected patients: a retrospective study of two centers in China. *PLoS One*. 2014;9(7):e101943. doi:10.1371/journal.pone.0101943.
- [30] Hamada S, Ichiyasu H, Inaba M, et al. Prognostic impact of pre-existing interstitial lung disease in non-HIV patients with *Pneumocystis pneumonia*. *ERJ Open Res*. April 2020;6(2). doi:10.1164/ajrccm-conference.2020.A108.
- [31] Hardak E, Neuberger A, Yigla M, et al. Outcome of *Pneumocystis jirovecii* pneumonia diagnosed by polymerase chain reaction in patients without human immunodeficiency virus infection. *Respirology*. May 2012;17(4):681–686. doi:10.1111/j.1440-1843.2012.02158.x.
- [32] Huang X, Weng L, Yi L, et al. [Clinical features of acute respiratory failure due to pneumocystis pneumonia in non-HIV immunocompromised patients]. *Zhonghua Yi Xue Za Zhi*. October 18, 2016;96(38):3057–3061. doi:10.3760/cma.j.issn.0376-2491.2016.38.005.
- [33] Ideguchi S, Yamamoto K, Hirayama T, et al. Diagnostic evaluation of serum (1, 3)- β -d-glucan levels using the Fungitec G-test MK kit for *Pneumocystis jirovecii* pneumonia (PCP) in non-HIV patients. *Med Mycol*. December 28, 2020, PMID: 33369648.
- [34] Inoue N, Fushimi K. Adjunctive corticosteroids decreased the risk of mortality of non-HIV pneumocystis pneumonia. *Int J Infect Dis*. February 2019;79:109–115. doi:10.1016/j.ijid.2018.12.001.
- [35] Issa N, Gabriel F, Baulier G, et al. Pneumocystosis and quantitative PCR. *Med Mal Infect*. October 2018;48(7):474–480. doi:10.1016/j.medmal.2018.04.396.
- [36] Jin F, Xie J, Wang HL. Lymphocyte subset analysis to evaluate the prognosis of HIV-negative patients with pneumocystis pneumonia. *BMC Infect Dis*. May 14, 2021;21(1):441. doi:10.1186/s12879-021-06124-5.
- [37] Kanj A, Samhouri B, Abdallah N, et al. Host factors and outcomes in hospitalizations for *Pneumocystis jirovecii* pneumonia in the United States. *Mayo Clin Proc*. February 2021;96(2):400–407. doi:10.1016/j.mayocp.2020.07.029.
- [38] Kato H, Samukawa S, Takahashi H, et al. Diagnosis and treatment of *Pneumocystis jirovecii* pneumonia in HIV-infected or non-HIV-infected patients—difficulties in diagnosis and adverse effects of trimethoprim-sulfamethoxazole. *J Infect Chemother*. November 2019;25(11):920–924. doi:10.1016/j.jiac.2019.06.007.

- [39] Kim T, Park SY, Lee HJ, et al. Assessment of cytomegalovirus and cell-mediated immunity for predicting outcomes in non-HIV-infected patients with *Pneumocystis jirovecii* pneumonia. *Medicine (Baltimore)*. July 2017;96(30):e7243. doi:10.1097/MD.0000000000007243.
- [40] Ko RE, Na SJ, Huh K, et al. Association of time-to-treatment with outcomes of *Pneumocystis pneumonia* with respiratory failure in HIV-negative patients. *Respir Res*. September 26, 2019;20(1):213. doi:10.1186/s12931-019-1188-6.
- [41] Kofteridis DP, Valachis A, Velegraki M, et al. Predisposing factors, clinical characteristics and outcome of *Pneumocystis jirovecii* pneumonia in HIV-negative patients. *J Infect Chemother*. July 2014;20(7):412–416. doi:10.1016/j.jiac.2014.03.003.
- [42] Korkmaz Ekren P, Töreyn ZN, Nahid P, et al. The association between cytomegalovirus co-infection with *Pneumocystis pneumonia* and mortality in immunocompromised non-HIV patients. *Clin Respir J*. November 2018;12(11):2590–2597. doi:10.1111/crj.12961.
- [43] Kosaka M, Ushiki A, Ikuyama Y, et al. A four-center retrospective study of the efficacy and toxicity of low-dose trimethoprim-sulfamethoxazole for the treatment of pneumocystis pneumonia in patients without HIV infection. *Antimicrob Agents Chemother*. December 2017;61(12):e01173-17. doi:10.1128/AAC.01173-17.
- [44] Kumagai S, Arita M, Koyama T, et al. Prognostic significance of crazy paving ground glass opacities in non-HIV *Pneumocystis jirovecii* pneumonia: an observational cohort study. *BMC Pulm Med*. February 21, 2019;19(1):47. doi:10.1186/s12890-019-0813-y.
- [45] Kunihiro Y, Tanaka N, Matsumoto T, et al. The usefulness of a diagnostic method combining high-resolution CT findings and serum markers for cytomegalovirus pneumonia and pneumocystis pneumonia in non-AIDS patients. *Acta Radiol*. July 2015;56(7):806–813. doi:10.1177/0284185114539320.
- [46] Lahmer T, da Costa CP, Held J, et al. Usefulness of 1,3 beta-D-Glucan detection in non-HIV immunocompromised mechanically ventilated critically ill patients with ARDS and suspected *Pneumocystis jirovecii* pneumonia. *Mycopathologia*. August 2017;182(7–8):701–708. doi:10.1007/s11046-017-0132-x.
- [47] Lee HY, Choi SH, Kim T, et al. Epidemiologic trends and clinical features of *Pneumocystis jirovecii* pneumonia in non-HIV patients in a tertiary-care hospital in Korea over a 15-year-period. *Jpn J Infect Dis*. July 24, 2019;72(4):270–273. doi:10.7883/yoken.JJID.2018.400.
- [48] Lemiale V, Debrumetz A, Delannoy A, et al. Adjunctive steroid in HIV-negative patients with severe *Pneumocystis pneumonia*. *Respir Res*. August 28, 2013;14(1):87. doi:10.1186/1465-9921-14-87.
- [49] Li LJ, Hsu SH, Wang CL, et al. Characteristics of viral pneumonia in non-HIV immunocompromised and immunocompetent patients: a retrospective cohort study. *BMC Infect Dis*. August 2021;21(1):767. doi:10.1186/s12879-021-06437-5.
- [50] Liu Y, Su L, Jiang SJ, et al. Risk factors for mortality from pneumocystis carinii pneumonia (PCP) in non-HIV patients: a meta-analysis. *Oncotarget*. August 29, 2017;8(35):59729–59739. doi:10.18632/oncotarget.19927.
- [51] Mansharamani NG, Garland R, Delaney D, et al. Management and outcome patterns for adult *Pneumocystis carinii* pneumonia, 1985 to 1995: comparison of HIV-associated cases to other immunocompromised states. *Chest*. September 2000;118(3):704–711. doi:10.1378/chest.118.3.704.
- [52] Matsumura Y, Shindo Y, Iinuma Y, et al. Clinical characteristics of *Pneumocystis pneumonia* in non-HIV patients and prognostic factors including microbiological genotypes. *BMC Infect Dis*. March 25, 2011;11(1):76. doi:10.1186/1471-2334-11-76.
- [53] Moon SM, Kim T, Sung H, et al. Outcomes of moderate-to-severe *Pneumocystis pneumonia* treated with adjunctive steroid in non-HIV-infected patients. *Antimicrob Agents Chemother*. October 2011;55(10):4613–4618. doi:10.1128/AAC.00669-11.
- [54] Mu XD, Jia P, Gao L, et al. Relationship between radiological stages and prognoses of pneumocystis pneumonia in non-AIDS immunocompromised patients. *Chin Med J (Engl)*. September 5, 2016;129(17):2020–2025. doi:10.4103/0366-6999.189068.
- [55] Mundo W, Morales-Shnaider L, Tewahade S, et al. Lower mortality associated with adjuvant corticosteroid therapy in non-HIV-infected patients with pneumocystis jirovecii pneumonia: a single-institution retrospective US cohort study. *Open Forum Infect Dis*. September 2020;7(9):ofaa354. doi:10.1093/ofid/ofaa354.
- [56] Nakamura H, Tateyama M, Tasato D, et al. Clinical utility of serum beta-D-glucan and KL-6 levels in *Pneumocystis jirovecii* pneumonia. *Intern Med*. 2009;48(4):195–202. doi:10.2169/internalmedicine.48.1680.
- [57] Nakashima K, Aoshima M, Nakashita T, et al. Low-dose trimethoprim-sulfamethoxazole treatment for pneumocystis pneumonia in non-human immunodeficiency virus-infected immunocompromised patients: a single-center retrospective observational cohort study. *J Microbiol Immunol Infect*. December 2018;51(6):810–820. doi:10.1016/j.jmii.2017.07.007.
- [58] Pereira-Díaz E, Moreno-Verdejo F, de la Horra C, et al. Changing trends in the epidemiology and risk factors of pneumocystis pneumonia in Spain. *Front Public Health*. 2019;7:275. doi:10.3389/fpubh.2019.00275.
- [59] Roblot F, Godet C, Le Moal G, et al. Analysis of underlying diseases and prognosis factors associated with *Pneumocystis carinii* pneumonia in immunocompromised HIV-negative patients. *Eur J Clin Microbiol Infect Dis*. July 2002;21(7):523–531. doi:10.1007/s10096-002-0758-5.

- [60] Roembke F, Heinzow HS, Gosseling T, et al. Clinical outcome and predictors of survival in patients with pneumocystis jirovecii pneumonia—results of a tertiary referral centre. *Clin Respir J*. January 2014;8(1):86–92. doi:10.1111/crj.12042.
- [61] Roux A, Canet E, Valade S, et al. Pneumocystis jirovecii pneumonia in patients with or without AIDS, France. *Emerg Infect Dis*. September 2014;20(9):1490–1497. doi:10.3201/eid2009.131668.
- [62] Sheikholeslami MF, Sadraei J, Farnia P, et al. Colonization of Pneumocystis jirovecii in Chronic Obstructive Pulmonary Disease (COPD) patients and the rate of Pneumocystis pneumonia in Iranian non-HIV(+) immunocompromised patients. *Iran J Microbiol*. December 2013;5(4):411–417.
- [63] Sun R, Lv D, Xiao M, et al. Diagnostic accuracy of the 1,3-beta-D-glucan test and lactate dehydrogenase for pneumocystis pneumonia in non-HIV patients. *Sci Rep*. April 29, 2021;11(1):9226. doi:10.1038/s41598-021-88729-z.
- [64] Tamai K, Tachikawa R, Tomii K, et al. Prognostic value of bronchoalveolar lavage in patients with non-HIV pneumocystis pneumonia. *Intern Med*. 2014;53(11):1113–1117. doi:10.2169/internalmedicine.53.0520.
- [65] Tang G, Tong S, Yuan X, et al. Using routine laboratory markers and immunological indicators for predicting Pneumocystis jirovecii pneumonia in immunocompromised patients. *Front Immunol*. 2021;12:652383. doi:10.3389/fimmu.2021.652383.
- [66] Tasaka S, Tokuda H, Sakai F, et al. Comparison of clinical and radiological features of pneumocystis pneumonia between malignancy cases and acquired immunodeficiency syndrome cases: a multicenter study. *Intern Med*. 2010;49(4):273–281. doi:10.2169/internalmedicine.49.2871.
- [67] Toper C, Rivaud E, Daniel C, et al. [Pneumocystis jirovecii pneumonia in non-HIV infected patients: a study of 41 cases]. *Rev Pneumol Clin*. September 2011;67(4):191–198. Pneumocystose pulmonaire chez des patients immunodéprimés non infectés par le VIH : étude de 41 cas. doi:10.1016/j.pneumo.2011.06.001.
- [68] Vogel MN, Brodoefel H, Hierl T, et al. Differences and similarities of cytomegalovirus and pneumocystis pneumonia in HIV-negative immunocompromised patients thin section CT morphology in the early phase of the disease. *Br J Radiol*. July 2007;80(955):516–523. doi:10.1259/bjr/39696316.
- [69] Weng L, Huang X, Chen L, et al. Prognostic factors for severe Pneumocystis jirovecii pneumonia of non-HIV patients in intensive care unit: a bicentric retrospective study. *BMC Infect Dis*. September 29, 2016;16(1):528. doi:10.1186/s12879-016-1855-x.
- [70] Wieruszewski PM, Barreto EF, Barreto JN, et al. Preadmission corticosteroid therapy and the risk of respiratory failure in adults without HIV presenting with pneumocystis pneumonia. *J Intensive Care Med*. December 2020;35(12):1465–1470. doi:10.1177/0885066619834242.
- [71] Wu HY, Wu KS, Huang YL, et al. Identifying predictors for bacterial and fungal coinfection on chest computed tomography in patients with Pneumocystis pneumonia. *J Microbiol Immunol Infect*. August 2021;54(4):701–709. doi:10.1016/j.jmii.2020.06.007.
- [72] Yu Q, Jia P, Su L, et al. Outcomes and prognostic factors of non-HIV patients with pneumocystis jirovecii pneumonia and pulmonary CMV co-infection: a retrospective cohort study. *BMC Infect Dis*. June 5, 2017;17(1):392. doi:10.1186/s12879-017-2492-8.
- [73] Zahar JR, Robin M, Azoulay E, et al. Pneumocystis carinii pneumonia in critically ill patients with malignancy: a descriptive study. *Clin Infect Dis*. October 15, 2002;35(8):929–934. doi:10.1086/342338.
- [74] Zalmanovich A, Ben-Ami R, Rahav G, et al. Healthcare-associated Pneumocystis jirovecii Pneumonia (PJP) Infection in HIV-negative adults: a multicenter study. *Isr Med Assoc J*. May 2021;23(5):312–317.
- [75] Wang D, Hu Y, Li T, Rong HM, Tong ZH. Diagnosis of Pneumocystis jirovecii pneumonia with serum cell-free DNA in non-HIV-infected immunocompromised patients. *Oncotarget*. September 22, 2017;8(42):71946–71953. doi:10.18632/oncotarget.18037.
- [76] Park JW, Curtis JR, Moon J, et al. Prophylactic effect of trimethoprim-sulfamethoxazole for pneumocystis pneumonia in patients with rheumatic diseases exposed to prolonged high-dose glucocorticoids. *Ann Rheum Dis*. May 2018;77(5):644–649. doi:10.1136/annrheumdis-2017-211796.
- [77] Truong J, Ashurst JV. Pneumocystis jirovecii pneumonia. In: *StatPearls*. Tampa, Florida, United States: Treasure Island (FL): StatPearls Publishing; January 21, 2023.