

Type of article: Original Research

Title: Comparative Therapeutic Strategies for Secondary Progressive Multiple Sclerosis: A Systematic Review and Network Meta-Analysis

Authors: Ye Yin¹, Jinmei Gao², Shahvaiz Magsi³, Danyu Zhao⁴

¹ Ultrasound Electrophysiology Center, Affiliated Hospital of Liaoning University of Traditional Chinese Medicine, Shenyang, China

² Department of Ultrasound, Shenyang Maternity and Child Health Hospital, Shenyang, Liaoning Province, China

³ Department of Neurology, West Virginia University, Clinton, United State

⁴ School of Basic Medical Sciences, Liaoning University of Traditional Chinese Medicine, Shenyang, China

Correspondence: Danyu Zhao

Address: No.79 Chongshan East Road, Huanggu District, Shenyang, Liaoning Province, China 110847

School of Basic Medical Sciences, Liaoning University of Traditional Chinese Medicine

Email: zhaody781107@163.com

Tel: +86 24 3120 7108

21
22
23
24
25
26
27
28
29

Clinical Question Box

What is the comparative effectiveness of disease-modifying therapies in adults with secondary progressive multiple sclerosis?

Moderate-certainty evidence suggests that siponimod may offer a favorable balance of efficacy and safety for delaying disability progression. Tolebrutinib appears promising but requires further investigation due to safety concerns. Interferon beta-1b remains a reasonable option, offering consistent benefits. Future head-to-head trials are needed to optimize treatment strategies for secondary progressive multiple sclerosis.

Abstract

Introduction: Secondary progressive multiple sclerosis (SPMS) represents a disabling stage of multiple sclerosis (MS) with limited treatment options. The evidence regarding the effectiveness of disease-modifying therapies (DMTs) in SPMS remains uncertain. This study evaluated the comparative efficacy and safety of DMTs in SPMS using a network meta-analysis (NMA).

Methods: A systematic search of PubMed, Embase, Web of Science, and Cochrane databases was conducted from January 2000 to May 2025 to identify randomized controlled trials (RCTs) involving adults with SPMS. The outcomes of interest were confirmed disability progression (CDP) and serious adverse events (SAEs). A frequentist random-effects NMA was performed to estimate odds ratios (ORs) and rank treatments using surface under the cumulative ranking (SUCRA) values.

Results: Seven RCTs involving 5,872 patients with SPMS were included in the final analysis. Approximately 60% of participants were women, with a mean age ranging from 39.9 to 48.9 years and a baseline Expanded Disability Status Scale (EDSS) score between 3.0 and 6.5. Tolebrutinib (OR 0.66, 95% confidence interval [CI] 0.50–0.87; SUCRA 0.713) and siponimod (OR 0.72, 95% CI 0.57–0.92; SUCRA 0.606) significantly reduced 6-month CDP compared with placebo. At 3 months, interferon beta-1b (OR 0.64, 95% CI 0.43–0.97; SUCRA 0.663), tolebrutinib (OR 0.73, 95% CI 0.56–0.96; SUCRA 0.554), and siponimod (OR 0.76, 95% CI 0.61–0.96; SUCRA 0.497) showed significant benefits. In terms of safety, tolebrutinib was associated with a higher risk of SAEs (OR 1.53, 95% CI 1.04–2.25), whereas the other agents had safety profiles comparable to placebo.

Conclusion: This NMA found that tolebrutinib, IFN- β -1b, and siponimod delay disability progression in SPMS, with IFN- β -1b and siponimod remaining the most practical options. The

52 conclusions should be interpreted cautiously given bias risk, moderate evidence certainty, limited
53 safety data, and clinical heterogeneity.

54 **Keywords:** secondary progressive multiple sclerosis, tolebrutinib, interferon beta-1b, siponimod,
55 network meta-analysis

Just Accepted

Introduction

MS is a chronic, immune-mediated demyelinating disease of the central nervous system, affecting nearly 1 million people in the United States and approximately 2.9 million people globally.¹ Among its clinical phenotypes, SPMS represents a particularly disabling stage that typically follows an initial relapsing-remitting phase (RRMS).² SPMS is characterized by gradual and irreversible neurological decline, with or without relapses, marking a shift from predominantly inflammatory to neurodegenerative pathology. Common symptoms include gait disturbances, spasticity, progressive motor impairment, and cognitive decline.³ The transition from RRMS to SPMS is often slow and difficult to diagnose early. It usually requires retrospective confirmation based on sustained progression. This diagnostic uncertainty hinders timely intervention, despite evidence suggesting that early detection and therapeutic escalation may improve long-term outcomes.⁴ Recent advances in imaging techniques, such as volumetric MRI and spinal cord analysis, along with emerging biomarkers like neurofilament light chain, are improving the early recognition of disease progression.^{5,6}

SPMS imposes a significant burden on both healthcare systems and individuals. A study conducted in Spanish neurology centers reported a mean annual cost of approximately €41,500 per patient with SPMS. Indirect costs—such as lost productivity, early retirement, and reduced leisure time—accounted for the majority of this amount. Direct medical expenses averaged €11,300, while non-medical costs, including home adaptations, totaled €8,800. In this cohort, over 90% of patients were unemployed, nearly 80% had taken early retirement, and caregivers lost more than 8 hours of work each week. Costs increased with the severity of disability, reaching €46,300 annually at an EDSS score of 6.5, compared to €34,850 at an EDSS score of 3 to 3.5, primarily

78 due to higher indirect costs.⁷ These findings highlight the substantial economic, personal, and
79 social impact of SPMS on both patients and healthcare systems.

80 Historically, SPMS, particularly its non-active form, was considered largely untreatable
81 due to the absence of overt inflammatory activity. However, recent approvals of DMTs for active
82 SPMS, characterized by ongoing relapses or new or enhancing MRI lesions, have transformed
83 clinical management.⁸ A major milestone was the approval of siponimod, a selective sphingosine
84 1-phosphate receptor modulator, following the EXPAND trial, which demonstrated significant
85 reductions in CDP, brain volume loss, and relapse rates.⁹ More recently, other oral agents such as
86 ozanimod and ponesimod have expanded the range of treatment options, each with distinct safety
87 and efficacy profiles.¹⁰ Ofatumumab, a CD20-targeting monoclonal antibody approved for PPMS
88 and RRMS, has also shown promise in active SPMS, particularly among patients transitioning
89 from RRMS.¹¹ Meanwhile, investigational therapies such as tolebrutinib, a Bruton's tyrosine
90 kinase inhibitor, are being studied in non-active SPMS. These therapies aim to target microglial-
91 driven neurodegeneration and intrathecal inflammation, which are key features of late-stage
92 disease.¹²

93 Ocrelizumab, cladribine, ofatumumab, and ponesimod are approved for the treatment of
94 RMS. Ocrelizumab, ofatumumab, and ponesimod received approval based on trials conducted in
95 RMS populations, whereas cladribine was approved following results from a study focused on
96 RRMS. Evidence supporting their efficacy in SPMS is limited and primarily derived from post
97 hoc analyses of small subgroups, typically comprising no more than 15% of trial participants. As
98 a result, the approval of these DMTs for RMS, including active SPMS, is largely based on the
99 assumption that relapse reduction observed in RRMS also applies to SPMS. However, the actual
100 effectiveness of these therapies in SPMS remains uncertain. This systematic review and NMA aim

101 to comprehensively evaluate the efficacy and safety of current and emerging treatments for SPMS,
102 using data exclusively from RCTs conducted in SPMS populations. The goal is to support
103 evidence-based decision-making in the management of this complex phase of the disease.

104

Just Accepted

Methods

Study Design and Registration

This systematic review and meta-analysis was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) 2020 guidelines. The review protocol was prospectively registered in the Open Science Framework database (OSF: osf.io/jhr6t).

Eligibility Criteria

We included studies that met the following criteria: (1) Population: adult patients aged 18 years or older diagnosed with SPMS; (2) Study Design: RCTs; (3) Intervention/Comparator: any pharmacological treatment compared with placebo or other active therapies; and (4) Outcomes: reporting at least one of the following: CDP (at 3 or 6 months) or SAEs. Studies were excluded if they (1) presented only subgroup analyses without new, relevant data; (2) had non-extractable or insufficient outcome data; (3) included fewer than 50% SPMS participants; (4) had a small sample size ($n < 50$) for SPMS participants; or (5) were non-randomized studies, conference abstracts, case reports, reviews, or animal studies.

Data bases

We searched PubMed, Embase, Web of Science, and the Cochrane Library from January 1, 2000, to May 31, 2025. In addition, we reviewed the reference lists of relevant articles and systematic reviews. A comprehensive search strategy was developed using the following search string: “((Randomized clinical trials) OR (RCT)) AND ((Secondary progressive multiple sclerosis) OR (SPMS))”.

Selection Process

Two independent reviewers (Reviewer Y.Y. and G.Z.) screened all retrieved titles and abstracts. Full texts of potentially relevant articles were assessed for eligibility. Discrepancies were resolved by discussion or a third reviewer (D.Z.). A standardized data extraction form was used by two reviewers to independently collect the following: study design, sample size, participant demographics, diagnostic criteria for SPMS, treatment arms, duration of follow-up, and reported outcomes (CDP and SAEs). Authors were contacted when essential data were missing or unclear.

Synthesis Methods

All statistical analyses were conducted using R version 4.4.1 (R Foundation for Statistical Computing, Vienna, Austria). Pairwise meta-analyses were performed using the “meta” package, and NMAs were conducted with the “netmeta” package, both based on frequentist methods. For binary outcomes (CDP and SAEs), ORs with 95% CIs were calculated. Consistency between direct and indirect evidence was assessed using node-splitting methods and global inconsistency tests. Treatment ranking was evaluated using P-scores, a frequentist analogue of the SUCRA curve, which estimates the probability of each treatment being the most effective or safest. Between-study heterogeneity was quantified using the I^2 statistic for pairwise comparisons and τ^2 for network models. A two-sided p-value < 0.05 was considered statistically significant.

Bias Assessment and Certainty of Evidence

The risk of bias for all included RCTs was assessed using the Cochrane Risk of Bias 2.0 tool. Two reviewers conducted the assessments independently, and any disagreements were resolved through consensus. Publication bias was evaluated using funnel plots and Egger’s regression test. Selective reporting was assessed by comparing published outcomes with study

148 protocols and registry entries, when available. The certainty of evidence for the primary outcomes
149 was assessed using the GRADE framework adapted for NMA. This evaluation considered five
150 domains: study limitations, inconsistency, indirectness, imprecision, and publication bias.

151

Just Accepted

Result

A total of 2,013 unique records were identified through systematic searches of the Cochrane, Embase, PubMed, and Web of Science databases. After initial screening, 188 full-text articles were assessed for eligibility. Of these, 181 were excluded for the following reasons: lack of a RCT design ($n = 78$), irrelevance to SPMS ($n = 57$), review articles ($n = 34$), unavailable data ($n = 10$), and small sample size ($n = 2$). Ultimately, seven studies met the inclusion criteria and were included in the final systematic review (Figure S1). These seven RCTs, comprising a total of 5,872 patients with SPMS, were included in the NMA (Table 1).¹³⁻¹⁹ The studies evaluated various DMTs, including low- and high-dose mitoxantrone (5 mg/m^2 and 10 mg/m^2), interferon beta-1a (IFN- β -1a), interferon beta-1b (IFN- β -1b) with both standard and adjusted dosages, natalizumab, siponimod, and tolebrutinib. Placebo arms served as comparators. The mean age of participants ranged from 39.9 to 48.9 years, with women accounting for approximately 60% of each study population. Most trials enrolled patients with EDSS scores between 3.0 and 6.5, indicating moderate to severe disability. Mitoxantrone was excluded from the NMA due to its known risks of cardiotoxicity and leukemia.

Six-Month CDP Outcomes

The network graph of the included studies is presented in Figure 1A. Two studies compared placebo with IFN- β -1b, while each of the other treatments was assessed in a single study. The direct comparisons are shown in Figure 2A. Among the interventions, low-dose mitoxantrone demonstrated the lowest OR for 6-month CDP at 0.28 (95% CI: 0.09–0.82), followed by tolebrutinib at 0.66 (95% CI: 0.50–0.87), IFN- β -1b at 0.68 (95% CI: 0.45–1.04), siponimod at 0.72 (95% CI: 0.57–0.92), IFN- β -1a at 0.85 (95% CI: 0.56–1.29), and natalizumab at 0.87 (95% CI: 0.62–1.22).

The results of the NMA are summarized in Table 2. Both tolebrutinib and siponimod demonstrated statistically significant superiority over placebo, with ORs consistent with those observed in direct comparisons. No other treatments showed superiority over one another. The relative efficacy of each treatment, based on SUCRA values, is illustrated in Figure S2: tolebrutinib (0.713), IFN- β -1b (0.645), siponimod (0.606), IFN- β -1a (0.403), IFN- β -1b with adjusted dosage (0.357), natalizumab (0.351), and placebo (0.176). There was no notable heterogeneity ($I^2 = 0\%$), and no evidence of publication bias was detected using Egger's test ($p = 0.90$).

Three-Month CDP Outcomes

The results for 3-month CDP outcomes were generally consistent with those observed at 6 months (Figure 1A). The direct comparisons are shown in Figure 2B. Low-dose and high-dose mitoxantrone demonstrated the lowest ORs for 3-month CDP, at 0.30 (95% CI: 0.10–0.90) and 0.63 (95% CI: 0.25–1.59), respectively. These were followed by IFN- β -1b at 0.64 (95% CI: 0.43–0.97), tolebrutinib at 0.73 (95% CI: 0.56–0.96), siponimod at 0.76 (95% CI: 0.61–0.96), adjusted-dose IFN- β -1b at 0.81 (95% CI: 0.40–1.64), IFN- β -1a at 0.96 (95% CI: 0.27–3.36), and natalizumab at 1.02 (95% CI: 0.47–2.23).

The results of the NMA are presented in Table 2. IFN- β -1b, tolebrutinib, and siponimod showed statistically significant superiority over placebo, with ORs consistent with those from the direct comparisons. No other agents demonstrated superiority over one another. The relative efficacy of each treatment, based on SUCRA values, is shown in Figure S3: IFN- β -1b (0.663), tolebrutinib (0.554), siponimod (0.497), adjusted-dose IFN- β -1b (0.428), IFN- β -1a (0.356), natalizumab (0.261), and placebo (0.196). There was no evidence of heterogeneity ($I^2 = 0\%$) and no publication bias was detected using Egger's test ($p = 0.58$).

The network graph of the enrolled studies is shown in Figure 1B. Only four agents reported data on SAEs. The results of the direct comparisons are presented in Figure 2C. IFN- β -1b showed an OR of 0.99 (95% CI: 0.63–1.58), natalizumab had an OR of 1.11 (95% CI: 0.81–1.53), siponimod showed an OR of 1.22 (95% CI: 0.92–1.61), and tolebrutinib had an OR of 1.53 (95% CI: 1.04–2.25).

The results of the NMA are presented in Table S1. Tolebrutinib was associated with a significantly higher risk of SAEs compared to placebo, with ORs consistent with those from the direct comparisons. No other agents showed significant differences from each other. The relative safety of each treatment, based on SUCRA values, is shown in Figure S4. The SUCRA values, rounded to three decimal places, were as follows: placebo (0.782), interferon beta-1a (IFN- β -1a) (0.709), natalizumab (0.548), siponimod (0.361), and tolebrutinib (0.101). Due to the small number of included studies, assessments of heterogeneity and publication bias using Egger's test are not considered reliable.

Risk of Bias and Certainty of Evidence

Risk of bias assessments (Figure S5) showed that four studies were rated as having some concerns, while the remaining three were judged to have a low risk of bias. The evaluated outcomes included CDP at 3 and 6 months. Overall, the risk of bias was considered to be of some concern. The certainty of evidence for the treatment rankings was rated as moderate, downgraded due to risk of bias.

Discussion

This NMA synthesized evidence from seven RCTs evaluating DMTs for SPMS. The results indicate that tolebrutinib, IFN- β -1b, and siponimod demonstrated significant benefits in delaying CDP at both 3 and 6 months compared to placebo. Notably, tolebrutinib was more effective than IFN- β -1b at 6 months, whereas IFN- β -1b showed greater benefit at 3 months. These findings are consistent with previous reports supporting the efficacy of IFN- β -1b and siponimod,²⁰ while also highlighting the emerging potential of tolebrutinib. In contrast, IFN- β -1a and natalizumab exhibited more modest effects, in line with earlier studies suggesting limited benefits in delaying disability progression.¹⁷ This review offers a comprehensive comparison of both established and newer agents by using an NMA framework to rank their relative effectiveness and safety.

Tolebrutinib, a next generation BTK inhibitor, demonstrated encouraging results with significant benefits at 3 and 6 months.²¹ Its dual mechanism of action, targeting peripheral immune cells and CNS resident microglia, offers a promising approach for SPMS, where compartmentalized inflammation drives progression. Nevertheless, concerns remain due to the higher risk of SAEs and the short follow up, which limits understanding of its long-term safety and efficacy. Real world evidence from observational cohorts, registries, and post marketing surveillance will be valuable in assessing durability of benefit, safety, and treatment persistence in broader patient populations. Furthermore, the inclusion of participants without active relapses in several trials may restrict generalizability. Despite these limitations, the findings highlight tolebrutinib's potential and underscore the need for longer term studies.

Regarding interferon agents, IFN- β -1b ranked slightly higher than IFN- β -1a, with lower ORs for CDP at both time points. This suggests a somewhat greater immunomodulatory effect, although the overall benefit remains modest and is unlikely to significantly influence long-term

outcomes. Similarly, natalizumab demonstrated limited benefit, with ORs close to unity, indicating no significant advantage in delaying disease progression. Mitoxantrone demonstrated the most pronounced short-term efficacy among the evaluated agents; however, its clinical utility is severely limited by a well-documented toxicity profile.²² Notably, the risks of cumulative dose-dependent cardiotoxicity and therapy-related secondary acute myeloid leukemia significantly outweigh its benefits in most settings, thereby precluding its routine use in long-term disease management.²³

From a clinical perspective, these findings suggest that siponimod remains the most practical treatment option due to its consistent efficacy and acceptable safety profile, as supported by current guidelines. Tolebrutinib appears promising, particularly for patients with active SPMS, but further evidence is needed to confirm its long-term benefits and safety. In contrast, interferon-based studies relied on older trial designs and outcome measures, which limit their applicability to contemporary clinical practice. Future research should prioritize head-to-head trials comparing tolebrutinib and siponimod in patients with active SPMS to clarify their relative effectiveness in slowing disease progression and preserving quality of life. Longer-term studies are also essential to assess sustained benefits, cognitive outcomes, and patient-reported measures. Additionally, improved standardization of trial criteria is necessary to enhance the generalizability of findings.

Several limitations of this review should be acknowledged. The risk of bias in the included studies raised some concern, and the certainty of evidence was moderate, which limits the strength of the conclusions. The small sample size and inclusion of multiple agents limited safety comparisons and precluded sensitivity analyses. While statistical heterogeneity was low, clinical heterogeneity, including differences in disease activity status, baseline EDSS, and trial durations, should be considered when interpreting the generalizability of these findings. Lastly, although no publication bias was detected, its presence cannot be entirely ruled out.

Conclusion

This systematic review and NMA demonstrated that tolebrutinib, IFN- β -1b, and siponimod are effective in delaying disability progression in SPMS. Although tolebrutinib shows promise, further research is needed to confirm its long-term efficacy and safety. IFN- β -1b and siponimod, with their consistent effectiveness and favorable safety profiles, remain the most practical treatment options. Future head-to-head trials will be critical to refining therapeutic strategies for SPMS.

271 **Acknowledgments:** None

272 **Funding Source:** None

273 **Author Contributions:** Y.Y. contributed to the study design and manuscript drafting. Y.Y. and
274 G.Z. performed the literature search, quality assessment, data extraction, and analysis. S.M. and
275 D.Z. contributed to data interpretation and revision. All authors have reviewed the manuscript and
276 agree with its content and data.

277 **Data Availability:** The corresponding author will provide the datasets used in this study upon
278 reasonable request.

279 **Ethical Statement:** Institutional Review Board approval was waived due to the nature of the meta-
280 analysis.

281 **Conflict of Interest:** The authors report no conflicts of interest related to this work.

282

- 284 1. Phillips CS, Kim J, Ganesh A, Stuifbergen AM. Impact of Active Versus Receptive Music
285 Interventions on Psychosocial and Neurological Outcomes in People with Multiple Sclerosis: A
286 Systematic Review. *J Integr Complement Med.* Jun 5 2025;doi:10.1089/jicm.2024.0588
- 287 2. Cree BAC, Arnold DL, Chataway J, et al. Secondary Progressive Multiple Sclerosis: New
288 Insights. *Neurology.* Aug 24 2021;97(8):378-388. doi:10.1212/wnl.00000000000012323
- 289 3. Ziemssen T, Bhan V, Chataway J, et al. Secondary Progressive Multiple Sclerosis: A
290 Review of Clinical Characteristics, Definition, Prognostic Tools, and Disease-Modifying
291 Therapies. *Neurol Neuroimmunol Neuroinflamm.* Jan
292 2023;10(1)doi:10.1212/nxi.000000000000200064
- 293 4. Rojas JI, Patrucco L, Alonso R, et al. Diagnostic uncertainty during the transition to
294 secondary progressive multiple sclerosis: Multicenter study in Argentina. *Mult Scler.* Apr
295 2021;27(4):579-584. doi:10.1177/1352458520924586
- 296 5. Bsteh G, Dal-Bianco A, Krajnc N, Berger T. Biomarkers of Progression Independent of
297 Relapse Activity-Can We Actually Measure It Yet? *Int J Mol Sci.* May 14
298 2025;26(10)doi:10.3390/ijms26104704
- 299 6. Rocca MA, Preziosa P, Barkhof F, et al. Current and future role of MRI in the diagnosis
300 and prognosis of multiple sclerosis. *Lancet Reg Health Eur.* Sep 2024;44:100978.
301 doi:10.1016/j.lanepe.2024.100978
- 302 7. Oreja-Guevara C, Meca-Lallana JE, Gómez-Estévez I, et al. Economic burden of
303 secondary progressive multiple sclerosis: DISCOVER study. *BMC Health Serv Res.* Apr 10
304 2025;25(1):525. doi:10.1186/s12913-025-12592-1
- 305 8. Mrochen A, Meuth SG, Pfeuffer S. Should we stay or should we go? Recent insights on
306 drug discontinuation in multiple sclerosis. *Neurol Res Pract.* Apr 21 2025;7(1):25.
307 doi:10.1186/s42466-025-00379-y
- 308 9. Hua LH, Bar-Or A, Cohan SL, et al. Effects of baseline age and disease duration on the
309 efficacy and safety of siponimod in patients with active SPMS: Post hoc analyses from the
310 EXPAND study. *Mult Scler Relat Disord.* Jul 2023;75:104766. doi:10.1016/j.msard.2023.104766
- 311 10. Dumitrescu L, Papathanasiou A, Coclitu C, et al. An update on the use of sphingosine 1-
312 phosphate receptor modulators for the treatment of relapsing multiple sclerosis. *Expert Opin*
313 *Pharmacother.* Mar 2023;24(4):495-509. doi:10.1080/14656566.2023.2178898
- 314 11. Bayas A, Christ M, Faissner S, et al. Disease-modifying therapies for relapsing/active
315 secondary progressive multiple sclerosis - a review of population-specific evidence from
316 randomized clinical trials. *Ther Adv Neurol Disord.* 2023;16:17562864221146836.
317 doi:10.1177/17562864221146836
- 318 12. Krämer J, Bar-Or A, Turner TJ, Wiendl H. Bruton tyrosine kinase inhibitors for multiple
319 sclerosis. *Nat Rev Neurol.* May 2023;19(5):289-304. doi:10.1038/s41582-023-00800-7
- 320 13. Hartung HP, Gonsette R, König N, et al. Mitoxantrone in progressive multiple sclerosis: a
321 placebo-controlled, double-blind, randomised, multicentre trial. *Lancet.* Dec 21-28
322 2002;360(9350):2018-25. doi:10.1016/s0140-6736(02)12023-x
- 323 14. Andersen O, Elovaara I, Färkkilä M, et al. Multicentre, randomised, double blind, placebo
324 controlled, phase III study of weekly, low dose, subcutaneous interferon beta-1a in secondary
325 progressive multiple sclerosis. *J Neurol Neurosurg Psychiatry.* May 2004;75(5):706-10.
326 doi:10.1136/jnnp.2003.010090

15. Panitch H, Miller A, Paty D, Weinshenker B. Interferon beta-1b in secondary progressive MS: results from a 3-year controlled study. *Neurology*. Nov 23 2004;63(10):1788-95. doi:10.1212/01.wnl.0000146958.77317.3e
16. Kappos L, Polman C, Pozzilli C, Thompson A, Beckmann K, Dahlke F. Final analysis of the European multicenter trial on IFNbeta-1b in secondary-progressive MS. *Neurology*. Dec 11 2001;57(11):1969-75. doi:10.1212/wnl.57.11.1969
17. Kapoor R, Ho PR, Campbell N, et al. Effect of natalizumab on disease progression in secondary progressive multiple sclerosis (ASCEND): a phase 3, randomised, double-blind, placebo-controlled trial with an open-label extension. *Lancet Neurol*. May 2018;17(5):405-415. doi:10.1016/s1474-4422(18)30069-3
18. Kappos L, Bar-Or A, Cree BAC, et al. Siponimod versus placebo in secondary progressive multiple sclerosis (EXPAND): a double-blind, randomised, phase 3 study. *Lancet*. Mar 31 2018;391(10127):1263-1273. doi:10.1016/s0140-6736(18)30475-6
19. Fox RJ, Bar-Or A, Traboulsee A, et al. Tolebrutinib in Nonrelapsing Secondary Progressive Multiple Sclerosis. *N Engl J Med*. May 15 2025;392(19):1883-1892. doi:10.1056/NEJMoa2415988
20. Cortesi PA, Antonazzo IC, Gasperini C, Nica M, Ritrovato D, Mantovani LG. Cost-effectiveness and budget impact analysis of siponimod in the treatment of secondary progressive multiple sclerosis in Italy. *PLoS One*. 2022;17(3):e0264123. doi:10.1371/journal.pone.0264123
21. Airas L, Bermel RA, Chitnis T, et al. A review of Bruton's tyrosine kinase inhibitors in multiple sclerosis. *Ther Adv Neurol Disord*. 2024;17:17562864241233041. doi:10.1177/17562864241233041
22. Brück W, Gold R, Lund BT, et al. Therapeutic decisions in multiple sclerosis: moving beyond efficacy. *JAMA Neurol*. Oct 2013;70(10):1315-24. doi:10.1001/jamaneurol.2013.3510
23. Shaikh AY, Suryadevara S, Tripathi A, et al. Mitoxantrone-Induced Cardiotoxicity in Acute Myeloid Leukemia-A Velocity Vector Imaging Analysis. *Echocardiography*. Aug 2016;33(8):1166-77. doi:10.1111/echo.13245

Table 1. Characteristics and Background of Enrolled Studies

Study	Cases	Ages (years)	Female (%)	Treatment	Route	Diagnosis	EDSS Score	Progression Criteria	FU
Hartung et al. 2002	188	39.9 (7.7)	98 (52.1%)	Mitoxantrone (L), Mitoxantrone (H), placebo	IV	Worsening RRMS or SPMS	3.0–6.0	≥1.0 EDSS point in past 18 months; stepwise or gradual progression; with/without relapses	2y
Andersen et al. 2004	364	45.8 (7.8)	224 (61.5%)	IFN-β-1a, placebo	SC	SPMS	< 7.0	≥1.0 EDSS point in past 4 years (0.5 if EDSS 6.0 or 6.5); ≥1 year MS duration; prior RRMS; ≥6 months worsening; with/without relapses	3y
Panitch et al. 2004	939	46.8 (0.5)	588 (62.6%)	IFN-β-1b (a), IFN-β-1b, placebo	SC	MS	3.0–6.5	≥1.0 EDSS point in past 2 years (0.5 if EDSS 6.5); ≥2 years MS duration; ≥1 relapse followed by ≥6 months progression	3y
Kappos et al. 2001	718	41.0 (7.2)	439 (61.1%)	IFN-β-1b, placebo	SC	SPMS	3.0–6.5	≥2 relapses or ≥1.0 EDSS increase in past 2 years	3y
Kapoor et al. 2018	887	47.3 (7.6)	550 (62.0%)	natalizumab, placebo	IV	SPMS	3.0–6.5	MSSS ≥4; progression unrelated to relapses in past year	3y
Kappos et al. 2018	1,645	48.1 (7.9)	992 (60.3%)	Siponimod, placebo	PO	SPMS	3.0–6.5	Prior RRMS; no relapse in past 3 months; EDSS progression in past 2 years	3y
Fox et al. 2025	1,131	48.9 (8.0)	696 (61.5%)	Tolebrutinib, placebo	PO	SPMS	3.0–6.5	Progression in past 12 months; no relapse in past 24 months	2y

Mitoxantrone (L): 5 mg/m²; Mitoxantrone (H): 10 mg/m²; FN-β-1b (a): 5 million unite/m²; IV: intravenous; SC: subcutaneous; PO: oral; y: year; SPMS: secondary progressive multiple sclerosis; EDSS: Expanded Disability Status Scale; MS: multiple sclerosis; MSSS: Multiple Sclerosis Severity Score; FU: follow-up

Table 2. Network Meta-Analysis of Various Treatments for Confirmed Disability Progression at 3 and 6 Months


Confirmed Disability Progression at six months						
Tolebrutinib						
0.97 [0.58; 1.60]	INF-β-1b					
0.91 [0.63; 1.32]	0.94 [0.58; 1.53]	Siponimod				
0.78 [0.47; 1.29]	0.80 [0.44; 1.46]	0.85 [0.52; 1.39]	INF-β-1a			
0.76 [0.49; 1.17]	0.78 [0.46; 1.34]	0.83 [0.55; 1.25]	0.97 [0.57; 1.67]	Natalizumab		
0.75 [0.42; 1.34]	0.77 [0.45; 1.32]	0.82 [0.46; 1.44]	0.96 [0.49; 1.87]	0.99 [0.54; 1.82]	INF-β-1b (a)	
0.66 [0.50; 0.87]	0.68 [0.45; 1.04]	0.72 [0.57; 0.92]	0.85 [0.56; 1.29]	0.87 [0.62; 1.22]	0.88 [0.53; 1.47]	Placebo
Confirmed Disability Progression at three months						
IFN-β-1b						
0.88 [0.54; 1.43]	Tolebrutinib					
0.84 [0.53; 1.35]	0.96 [0.68; 1.36]	Siponimod				
0.79 [0.38; 1.63]	0.90 [0.43; 1.90]	0.94 [0.45; 1.96]	IFN-β-1b (a)			
0.67 [0.18; 2.53]	0.77 [0.21; 2.77]	0.80 [0.22; 2.86]	0.85 [0.20; 3.58]	IFN-β-1a		
0.63 [0.26; 1.52]	0.72 [0.31; 1.64]	0.75 [0.33; 1.68]	0.80 [0.28; 2.27]	0.94 [0.21; 4.11]	Natalizumab	
0.64 [0.43; 0.97]	0.73 [0.56; 0.96]	0.76 [0.61; 0.96]	0.81 [0.40; 1.64]	0.96 [0.27; 3.36]	1.02 [0.47; 2.23]	placebo

FN-β-1b (a): 5 million unite/m² IV

Table 3. Network Meta-Analysis of Various Treatments for Serious Adverse Events


Placebo				
1.01 [0.63; 1.59]	INF-β-1a			
0.90 [0.65; 1.24]	0.90 [0.51; 1.57]	Natalizumab		
0.82 [0.62; 1.09]	0.82 [0.48; 1.40]	0.91 [0.60; 1.40]	Siponimod	
0.65 [0.44; 0.96]	0.65 [0.36; 1.19]	0.73 [0.44; 1.20]	0.80 [0.49; 1.29]	Tofacitinib

Figure 1. Network graph of enrolled studies



A: Studies CDP at 3 and 6 months; B: Studies on serious adverse events.

Figure 2. Direct Comparison of Various Treatments



A: Direct Comparison of CDP at 6 Months; B: Direct CDP Progression at 3 Months; C: Direct Comparison of the Risk of SAEs.